

GEM4 Summer School OpenCourseWare

<http://gem4.educommons.net/>

<http://www.gem4.org/>

Lecture: “Thermal Forces and Brownian Motion” by Ju Li.

Given August 11, 2006 during the GEM4 session at MIT in Cambridge, MA.

Please use the following citation format:

Li, Ju. “Thermal Forces and Brownian Motion.” Lecture, GEM4 session at MIT, Cambridge, MA, August 11, 2006.

<http://gem4.educommons.net/> (accessed MM DD, YYYY). License: Creative Commons Attribution-Noncommercial-Share Alike.

Note: Please use the actual date you accessed this material in your citation.

Thermal Forces and Brownian Motion

Ju Li



GEM4 Summer School 2006

Cell and Molecular Mechanics in BioMedicine

August 7–18, 2006, MIT, Cambridge, MA, USA

Outline

- Meaning of the Central Limit Theorem
- Diffusion vs Langevin equation descriptions
(average vs individual)
 - Diffusion coefficient and fluctuation-dissipation theorem

Central Limit Theorem

$$Y = X_1 + X_2 + \dots + X_N$$

X_1, X_2, \dots, X_N are random variables

$$E[Y] = E[X_1] + E[X_2] + \dots + E[X_N]$$

If X_1, X_2, \dots, X_N are *independent* random variables:

$$\text{var}[Y] = \text{var}[X_1] + \text{var}[X_2] + \dots + \text{var}[X_N]$$

Note: $\text{var}[X] = \sigma^2_X \equiv E[(X - E[X])^2]$

If X_1, X_2, \dots, X_N are *independent* random variables sampled from the *same* distribution:

$$E[Y] = NE[X]$$



$$\text{var}[Y] = N \text{var}[X_1] = N\sigma^2_X$$

Average of the sum: $y \equiv Y/N$

$$E[y] = E[X], \quad \text{var}[y] = \text{var}[Y]/N^2 = \sigma^2_X / N$$


Law of large numbers: as N gets large, the average of the sum becomes more and more deterministic, with variance σ^2_X / N .

X_1, X_2, \dots, X_N may be sampled from



We know the probability distribution of Y is shifting ($NE[X]$), as well as getting fat ($N\sigma^2_X$). But how about its *shape* ?

The central limit theorem says that *irrespective* of the shape of X ,



Why Gaussian ?

$$\rho(Y) \xrightarrow{\text{large } N} \frac{1}{\sqrt{2\pi N\sigma_X^2}} \exp\left(\frac{(Y - NE[X])^2}{2N\sigma_X^2}\right)$$

Gaussian is special

(Maxwellian velocity distribution, etc).

While proof is involved,
here we note that Gaussian is an invariant
shape (attractor in shape space) in the
mathematical operation of convolution.

Diffusion Equation in 1D

$$\partial_t \rho = -\partial_x \left(\overbrace{-D \partial_x \rho}^J \right) = D \partial_x^2 \rho$$

Random walker view of diffusion: imagine

(a) We release the walker at $x=0$ at $t=0$,

(b) Walker makes a move of $\pm a$, with equal probability, every $\Delta t=1/v$ from then on.

Mathematically, we say $\rho(x, t=0) = \delta(x)$.

$N = \frac{t}{\Delta t} = vt$ independent random steps

$$\text{Then, } x(t) = \overbrace{\Delta x_1 + \Delta x_2 + \dots + \Delta x_{t/\Delta t}}$$

When $N=vt \gg 1$,
the central limit theorem applies:

$$E[x(t)] = 0, \quad \text{var}[x(t)] = vt \text{ var}[\Delta x] = vta^2$$

So we can directly write down $\rho(x(t))$ as

$$\rho_G(x, t) = \frac{1}{\sqrt{2\pi va^2 t}} \exp\left(-\frac{x^2}{2va^2 t}\right)$$

It is the probability of finding the walker at x
at time t , knowing he was at 0 at time 0.

By plugging in, we can directly verify

$\rho_G(x, t)$ satisfies


$$\partial_t \rho = D \partial_x^2 \rho, \quad \rho(x, 0) = \delta(x).$$

with macroscopic D identified as $\frac{va^2}{2}$.

$$\rho_G(x, t) = \frac{1}{\sqrt{2\pi(2Dt)}} \exp\left(-\frac{x^2}{2(2Dt)}\right)$$

is called Green's function solution
to diffusion equation.

Brownian Motion



Courtesy of Microscopy-UK. Used with permission.

Fat droplets suspended in milk (from Dave Walker).

The droplets range in size from about 0.5 to 3 μm .

viscous oil



Stokes' law:

$$F = -6\pi r\eta v = -\lambda v$$

$$m\dot{v} = F = -\lambda v, \quad v(t=0) = v_0$$

$$\rightarrow v(t) = v_0 e^{-\frac{\lambda}{m}t}$$

Einstein's Explanation of Brownian Motion

Also, equi-partition theorem: $\left\langle \frac{mv^2}{2} \right\rangle = \frac{k_B T}{2}$

In addition to dissipative force, there must be another, stimulative force.

$$m\dot{v} = F_{\text{dissipative}} + F_{\text{stimulative/fluctuation}} = -\lambda v + F_{\text{fluc}}(t)$$

$$\langle F_{\text{fluc}}(t) \rangle = 0$$

$$\langle F_{\text{fluc}}(t) F_{\text{fluc}}(t') \rangle = b(t - t')$$

If $b(t - t') = B\delta(t - t')$: white noise

Exact Green's function solution of $v(t)$:

$$v(t) = \frac{1}{m} \int_{-\infty}^t dt' F_{\text{fluc}}(t') e^{-\frac{\lambda}{m}(t-t')}$$

$$\langle v(t)v(\tilde{t}) \rangle$$

$$= \frac{1}{m^2} \left\langle \int_{-\infty}^t dt' F_{\text{fluc}}(t') e^{-\frac{\lambda}{m}(t-t')} \int_{-\infty}^{\tilde{t}} d\tilde{t}' F_{\text{fluc}}(\tilde{t}') e^{-\frac{\lambda}{m}(\tilde{t}-\tilde{t}')} \right\rangle$$

$$= \frac{1}{m^2} \int_{-\infty}^t dt' e^{-\frac{\lambda}{m}(t-t')} \int_{-\infty}^{\tilde{t}} d\tilde{t}' e^{-\frac{\lambda}{m}(\tilde{t}-\tilde{t}')} \langle F_{\text{fluc}}(t') F_{\text{fluc}}(\tilde{t}') \rangle$$

$$= \frac{1}{m^2} \int_{-\infty}^t dt' e^{-\frac{\lambda}{m}(t-t')} \int_{-\infty}^{\tilde{t}} d\tilde{t}' e^{-\frac{\lambda}{m}(\tilde{t}-\tilde{t}')} B \delta(t' - \tilde{t}')$$

$$= \frac{1}{m^2} \int_{-\infty}^t dt' e^{-\frac{\lambda}{m}(t-t')} H(\tilde{t} - t') e^{-\frac{\lambda}{m}(\tilde{t}-t')} B$$

$H(x)$ is Heaviside step function:

$$= \frac{B}{2m\lambda} e^{-\frac{\lambda}{m}|t-\tilde{t}|} \quad H(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x \leq 0 \end{cases}$$

In particular: $\langle v(t)v(t) \rangle = \frac{B}{2m\lambda}$

However, from equilibrium statistical mechanics:
equi-partition theorem:

$$m \langle v(t)v(t) \rangle = k_B T$$

$$\rightarrow \frac{B}{2\lambda} = k_B T$$

The ratio between *square* of stimulative force
and dissipative force is fixed, $\propto T$

$$\langle v(t)v(\tilde{t}) \rangle = \frac{k_B T}{m} e^{-\frac{\lambda}{m}|t-\tilde{t}|}$$

Previously, from the Gaussian solution to

$$\partial_t \rho = D \partial_x^2 \rho, \quad \rho(x, 0) = \delta(x) :$$

$$\rho_G(x, t) = \frac{1}{\sqrt{2\pi(2Dt)}} \exp\left(-\frac{x^2}{2(2Dt)}\right)$$

we know if the particle is released at $x = 0$ at $t = 0$:

$$\langle x(t)x(t) \rangle = 2Dt$$

$$x(t) = 0 + \int_0^t dt' v(t'), \quad \dot{x}(t) = v(t)$$

$$\begin{aligned} \frac{d}{dt} \langle x(t)x(t) \rangle &= 2 \langle x(t)\dot{x}(t) \rangle = 2 \langle x(t)v(t) \rangle \\ &= \frac{d}{dt} (2Dt) = 2D \end{aligned}$$

$$\begin{aligned} D &= \langle x(t)v(t) \rangle = \left\langle \left(\int_0^t dt' v(t') \right) v(t) \right\rangle \\ &= \int_0^t dt' \langle v(t')v(t) \rangle \\ &= \int_0^t dt' \langle v(t')v(0) \rangle \end{aligned}$$

Velocity auto-correlation function: $g(t) \equiv \langle v(t)v(0) \rangle$

Actually, the onset of macroscopic diffusion

$(\partial_t \rho = D \partial_x^2 \rho)$ is only valid only when

$t \gg$ intrinsic timescale of $g(t) \propto \frac{m}{\lambda}$

(Same as central limit theorem in random walk)

So the correct formula is

$$D = \int_0^{\infty} dt' \langle v(t') v(0) \rangle$$

The above is one of the
fluctuation-dissipation theorems.

Thermal conductivity: $\kappa = \frac{1}{\Omega k_B T^2} \int_0^\infty \langle J_q(t) J_q(0) \rangle dt$

Electrical conductivity: $\sigma = \frac{1}{\Omega k_B T} \int_0^\infty \langle J(t) J(0) \rangle dt$

Shear viscosity: $\eta = \frac{\Omega}{k_B T} \int_0^\infty \langle \tau_{xy}(t) \tau_{xy}(0) \rangle dt$

Fluctuation-dissipation theorem (Green-Kubo formula) is one of the most elegant and significant results of statistical mechanics. It relates *transport properties* (system behavior if linearly perturbed from equilibrium) to the time-correlation of *equilibrium fluctuations*.

Coming back to diffusion (mass transport):

$$\langle v(t)v(\tilde{t}) \rangle = \frac{k_{\text{B}}T}{m} e^{-\frac{\lambda}{m}|t-\tilde{t}|}$$

$$\text{So } D = \int_0^\infty dt' \langle v(t')v(0) \rangle = \frac{k_{\text{B}}T}{\lambda}.$$

$\frac{1}{\lambda}$ is actually the mobility of the particle, when driven by external (non-thermal) force.

$\frac{D}{1/\lambda} = k_{\text{B}}T$ is called the Einstein relation,
first derived in 1905.

References

Kubo, Toda & Hashitume,
Statistical Physics II: Nonequilibrium Statistical Mechanics
(Springer-Verlag, New York, 1992).

Zwanzig, Nonequilibrium Statistical Mechanics
(Oxford University Press, Oxford, 2001).

van Kampen, Stochastic processes in physics and chemistry,
rev. and enl. ed. (North-Holland, Amsterdam, 1992).

Reichl, A modern course in statistical physics
(Wiley, New York, 1998).