


# Biochemistry Unit Exam

## Question 1


a) An example of a structural representation is shown in the adjacent box.

Draw a structural representation of the amino acid, Aspartic acid, which has the side chain of:  $\text{CH}_2\text{COOH}$ .




b) The molecule drawn above is a monomer found in \_\_\_\_\_.

c) Below is a pentose found in RNA. On the diagram circle the hydroxyl that differs between ribonucleotides and deoxyribonucleotides.



d) You find an organism that lives in thermal pools with temperatures as high as  $160^\circ\text{F}$ . You find that many of this organism's proteins have a high percentage of cysteine. Briefly explain why this might be the case.


e) To the right is an iconic view of a phospholipid.



- List what atoms are found in a phospholipid molecule represented by the boxed region of this schematic. Describe the properties associated with this region.
- List what atoms are found in a phospholipid molecule represented by the shaded region of this schematic. Describe the properties associated with this region.

**Question 1, continued**


f)



- a) Box the part that is added to a growing chain of nucleic acid.
- b) Star the atom(s) that can form a hydrogen bond with the complementary nitrogenous base.
- c) Circle the part of the molecule that decreases the stability of RNA as compared to DNA.
- d) Draw an arrow to the part of this molecule that you would modify to prevent further elongation. Indicate what change you would make next to the arrow drawn.

**Question 2**

Shown below is the structure of Drug 1, that binds to the *E. coli* ribosome. Indicate whether each circled region could possibly form ionic bonds, hydrogen bonds, or hydrophobic interactions with another molecule by filling out the table below. Fill in the table with “yes” if that type of bond is possible, “no” if it is not.




a)

<b>Part</b>	Could this part form <b>ionic bonds</b>	Could this part form <b>hydrogen bonds</b>	Could this part be involved in a <b>hydrophobic interaction</b>
(i)			
(ii)			
(iii)			
(iv)			

**Question 2, continued,**

A representation of the drug bound to the *E. coli* ribosome is shown below. Three amino acids of the ribosome that are important for binding are shown. If this drug can bind to different bacterial ribosomes, it may be useful as an antibiotic. The ability of this drug to bind to the ribosomes of two different species is examined. The drug binds to the ribosome of species 1, but does **not** bind to the ribosome of species 2.



For your information:

Amino acid	Side chain
Ile	CHCH <sub>3</sub> CH <sub>2</sub> CH <sub>3</sub>
Val	CHCH <sub>3</sub> CH <sub>3</sub>
Asn	CH <sub>2</sub> CONH <sub>2</sub>
Glu	CH <sub>2</sub> CH <sub>2</sub> COO <sup>-</sup>
Lys	(CH <sub>2</sub> ) <sub>4</sub> NH <sub>3</sub> <sup>+</sup>
Ser	CH <sub>2</sub> OH
Leu	CH <sub>2</sub> CHCH <sub>3</sub> CH <sub>3</sub>
Arg	(CH <sub>2</sub> ) <sub>3</sub> NHC(NH <sub>2</sub> ) <sub>2</sub> <sup>+</sup>


b) The table shows the differences between the ribosomes. Given this information, complete the table indicating which would be from species 1 (drug binds) and which would be from species 2 (drug does not bind).

Ribosome from:	Amino acid found at		
	101	231	312
<i>E. coli</i>	Lys	Asn	Leu
Species _____	Glu	Ser	Val
Species _____	Arg	Glu	Ile

c) What specific amino acid or amino acids prevents the ribosome from species 2 from binding Drug 1?


d) In terms of the specific non-covalent interactions, explain why the ribosome from species 2 will NOT bind Drug 1, but the ribosomes from *E. coli* and species 1 will.

### Question 3



a) The energy profile for the reaction 1 is drawn on the axes below.

- Label A, B, and  $\Delta G'$  on the graph below.
- What is the value for the energy of activation?



Compare reaction 1 (above) to reactions 2 and 3 (below).



b) Which of these reactions, 1, 2, or 3 is most likely to proceed in the forward direction in the absence of an enzyme? If you do not have enough information to answer this question, write "Can't tell" below. Explain your answer.

c) When an appropriate enzyme is added to each of these reactions, the rate of the reaction increases. Which reaction will proceed the fastest? If you do not have enough information to answer this question, write "Can't tell" below. Explain your answer.

d) Assume that you have a reaction  $X \rightleftharpoons Y$  where  $\Delta G = 0$  and the energy of activation = +0.6 kcal/mole. In a cell, you find that the reaction proceeds almost exclusively in the forward direction. Explain why this might be the case.

### Question 3, continued

Enzyme 1 carries out the reaction:  $W \rightarrow X + Y$ . There are three regions where enzyme 1 interacts with substrate W (another protein) to facilitate the conversion of W into X and Y.

Region 1 is stabilized by hydrophobic interactions between the side chains of the amino acids found on the enzyme and substrate.

Region 2 is stabilized by hydrogen bonds between the side chains of the amino acids found on the enzyme and substrate.

Region 3 is stabilized by ionic bonds between the side chains of the amino acids found on the enzyme and substrate.

e) You have the following pairs of amino acids. For each pair given, list in which region (1, 2, or 3) this pair would act to stabilize the interaction of enzyme 1 with substrate A. If the pair could be found in more than one region, list all that apply. If the pair would not be found in any region write in the word **NONE**.

pair	Amino acid on enzyme	Amino acid on substrate	In what region or regions would the pair be found?
1	alanine	leucine	
2	arginine	isoleucine	
3	serine	methionine	
4	threonine	serine	
5	lysine	arginine	
6	glutamine	asparagine	
7	proline	valine	
8	aspartic acid	lysine	

### Question 4

You inoculate two test tubes with the same amount of identical growth medium and with the same number of identical yeast cells and grow these cells under identical conditions except for the presence or absence of oxygen.

a) After 12 hours, all of the glucose in each culture has been consumed. You determine the number of total cells found in each culture and find that one culture has more cells than the other.

- Which culture would have the greater cell density, the one grown aerobically or the one grown anaerobically?
- Explain why the culture you chose above can make more cells with the same amount of glucose than the other culture.

b) The cells of both cultures convert glucose to pyruvate via glycolysis, and then further metabolize pyruvate.

- Are the cells of both cultures able to obtain the same amount of ATP from **glycolysis**? Explain.
- Under anaerobic conditions, the carbon from pyruvate will ultimately be found in which molecule?
- Under aerobic conditions, the carbon from pyruvate will ultimately be found in which molecule?

**Question 4, continued**

c) Under aerobic conditions, when glucose is metabolized some of the energy is used to reduce  $\text{NAD}^+$  to  $\text{NADH} + \text{H}^+$ . In the mitochondria,  $\text{NADH}$  donates its electrons to  $\text{NADH-Q}$  reductase, and from there the electrons move through the electron transport chain.

- Explain why this process requires aerobic conditions.
- Briefly describe how the transfer of electrons from one protein to another in the electron transport chain results in the production of ATP.

d) In oxygenic photosynthesis, electrons from the chlorophyll found in photosystem II are donated to the primary electron carrier. How are the donated electrons replaced?

e) Anoxygenic photosynthetic organisms make ATP using an electron transport chain, but do **not** produce  $\text{O}_2$  as a waste product. What is a common source of electrons for the conversion of  $\text{NADP}^+$  to  $\text{NADPH}$  for in this type of organism?

*Many highly reduced compounds whose redox potential will allow transfer of electrons to  $\text{NADP}^+$  could serve as a source of electron. One common compound is  $\text{H}_2\text{S}$ .*

f) Circle **all** of the metabolic processes that occur in **organisms** that perform oxygenic photosynthesis.

Glycolysis	Citric acid Cycle	Oxidative phosphorylation
Calvin Cycle	Cyclic photophosphorylation	Non-cyclic photophosphorylation.

g) Both respiration and photosynthesis are evolutionarily conserved. Based upon the current understanding of how life on earth evolved, which of these processes likely evolved first? Justify your answer in the space below.

MIT OpenCourseWare  
<http://ocw.mit.edu>

7.01SC Fundamentals of Biology  
Fall 2011

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.