

MIT OpenCourseWare  
<http://ocw.mit.edu>

5.111 Principles of Chemical Science  
Fall 2008

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

## 5.111 Lecture 27

Transition Metals                      Topics: Formation of coordination complexes, coordination number, coordination complex notation, structures of coordination complexes, chelate effect, isomers, d-electron counting, and d-orbitals.

Chapter 16 p. 669-681 (p. 620-631 in 3<sup>rd</sup> ed).

From Wednesday's handout

Now the answer to the biochemical question

How is vitamin B<sub>12</sub> reduced in the body? Vitamin B<sub>12</sub> is reduced by a protein called flavodoxin.

$E^\circ$  for vitamin B<sub>12</sub> is -0.526 V

$E^\circ$  for flavodoxin is -0.230 V

Is the reduction of vitamin B<sub>12</sub> by flavodoxin spontaneous?

$$\begin{aligned}\Delta E^\circ(\text{cell}) &= E^\circ(\text{reduction}) - E^\circ(\text{oxidation}) \\ &= E^\circ(\text{vitamin B}_{12}) - E^\circ(\text{flavodoxin}) \\ &= -0.526 \text{ V} - (-0.230 \text{ V}) = -0.296 \text{ V}\end{aligned}$$

$$\Delta G^\circ = -n\mathfrak{F}\Delta E^\circ = -(1)(96485 \text{ Cmol}^{-1})(-0.296 \text{ V}) = +28.6 \text{ kJ/mol}$$

Vitamin B<sub>12</sub> is a better reducing agent than flavodoxin. Vitamin B<sub>12</sub> should reduce flavodoxin not the other way around. So why don't we all have heart disease and megaloblastic anemia?

Answer: S-adenosylmethionine provides the energy to drive the reaction. The  $\Delta G^\circ$  for the cleavage of S-adenosylmethionine is -37.6 kJ/mol

Today's material

21 Sc	22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn
39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd
		74 W					78 Pt	79 Au	80 Hg

Developed from Lippard & Berg 1994

d-block metals naturally occurring in biology – V, Cr, Mn, Fe, Co, Ni, Cu, Zn, Mo, Cd, W.

d-block metals used as probes of biological systems and/or drugs include Cr, Co, Y, Tc, Ag, Cd, Pt, Au, Hg.

Roles of metals in biology include  
global cycling of nitrogen, carbon, hydrogen  
 radical reactions  
 biosynthesis of vitamins  
 biosynthesis of deoxynucleotides  
 etc


### Formation of coordination complexes

A key feature of transition metals is their ability to form complexes with small molecules and ions.

Positive metals ions can attract electron density, usually a lone pair of electrons from another atom or molecule to form a coordination complex.

Donor atoms are called ligands (Lewis \_\_\_\_\_ –typically \_\_\_\_\_ one lone pair of electrons)


Examples of ligands:



Acceptor atoms are transition metals (Lewis acid –accept lone pair electrons)

Examples of transition metals: Ti, Cr, Mn, Fe, Co, Ni, Zn, Ir, Pt, etc



Coordination complexes = metals surrounded by ligands. Example:



Coordination number (CN) is the number of ligands bonded to the metal ion. Here CN = 6. Six ligands comprise the primary coordination sphere.


Typical CN's range from 2-12. Six is the most common.

### Coordination Complex Notation



$\text{NH}_3$  within bracket is bound to Co, Cl outside bracket is a counter ion.

### Structures of coordination complexes (M = Metal, L = Ligand)



### Chelate Effect in Coordination Complexes

Ligands that bind a metal at one site are called unidentate or monodentate (one tooth).

Ligands that have two or more points of attachment to the metal are called chelating ligands and the coordination complexes are called chelates (greek for claws).

Bidentate – two points of attachment

Tridentate –

Tetradentate -

ex. corrin ring of  $\text{B}_{12}$

Hexadentate –


ex. EDTA

Metal chelates are unusual stable. This is partly due to the favorable entropic factor accompanying release of non-chelating ligands (usually  $\text{H}_2\text{O}$ ) from the coordination sphere.


Examples

### 1. Vitamin B<sub>12</sub>.


Cobalt is coordinated by a planar tetradentate ligand (corrin ring system). It is also coordinated by an upper axial ligand (5'-deoxyadenosine) and a lower axial ligand (dimethylbenzimidazole).



### 2. Ethylenediamine tetraacetic acid (EDTA).



Free EDTA



EDTA in complex with metal (M)


Binding of EDTA is entropically favorable. Six molecules of H<sub>2</sub>O are released for every 1 molecule of EDTA bound.

### Uses


## Geometric Isomers

Geometric isomers can have vastly different properties.

$[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$  has two geometric isomers




cisplatin



transplatin

Optical isomers (enantiomers) are non-superimposable mirror images of each other.

A complex that is not identical to its mirror image is also called a chiral complex. Chiral molecules have different properties in chiral environments (such as a human body).



## d-Electron Counting in Coordination Complexes

d-electron count of metal = group number (periodic table) - oxidation number of metal

1. find oxidation number:

For Co in  $[\text{Co}(\text{NH}_3)_6]^{3+}$

$\text{NH}_3$  is neutral, so Co must be +3

2. d-count is  $9 - 3 = 6$   $d^6$

## Practice with d-counts


Complexes

Oxidation number of metal


d-count

 $[\text{Ni}(\text{CO})_4]$  $[\text{Co}(\text{H}_2\text{O})_2(\text{NH}_3)\text{Cl}_3]^-$ d OrbitalsThere are five d orbitals:  $d_{xy}$ ,  $d_{xz}$ ,  $d_{yz}$ ,  $d_{x^2-y^2}$ ,  $d_z^2$ .


You need to be able to draw their shapes.




$d_z^2$  has maximum amplitude along z and doughnut in xy plane




$d_{x^2-y^2}$  has maximum amplitude along x and y axes.



$d_{yz}$  has maximum amplitude  $45^\circ$  to y and z axes



$d_{xz}$  has maximum amplitude  $45^\circ$  to x and z axes



$d_{xy}$  has maximum amplitude  $45^\circ$  to x and y axes