

Chapter 11
Nucleophilic Substitution (S_N1/S_N2)
Elimination (E1/E2)

Suggested Problems for Elimination:

11.14, 11.17-19, 11.20, 11.34, 11.40, 11.45-46, 11.49-50, 11.56, 11.59, 11.65, 11.68

Substitution Reactions (S_N2 versus S_N1)

S_N2

S_N1


	S_N2	S_N1
Mechanism:	Concerted	Two Steps (Look for carbocation rearrangements.)
Rate Equation:	Rate = $k_r[R-X][Nuc]$	Rate = $k_r[R-X]$
Stereochemistry:	Stereospecific (inversion)	Loss of Stereochemistry
Substrate:	Sterics: (methyl > 1° > 2°) No S_N2 with 3°! Accessible to Nu: (benzylic > allylic > 1°)	Cation Stability (benzylic > allylic > 3° > 2° >> 1°) No 1° or methyl R^+ without extra resonance stabilization!
Nucleophile:	Strong/Moderate Required strong: RS^- , I^- , R_2N^- , R_2NH , RO^- , CN^- moderate: RSH , Br^- , RCO_2^-	Not Important
Leaving Group:	Moderately Important ($-OTf \gg -OTs @ -OMs \gg -I > -Br > -Cl$)	Very Important ($-OTf \gg -OTs @ -OMs \gg -I > -Br > -Cl$)
Solvent:	Polar Aprotic	Polar Protic

Elimination Reactions: E2 versus E1


	E2	E1
Mechanism:	Concerted	Two Steps (Look for carbocation rearrangements.)
Rate Equation:	Rate = $k_r[R-X][Base]$	Rate = $k_r[R-X]$
Stereochemistry:	Stereospecific (antiperiplanar TS)	Not Stereospecific
Substrate:	Alkene Stability (3° > 2° > 1°)	Cation Stability (benzylic > allylic > 3° > 2° >> 1°)
Base:	Strong Base Required (RO ⁻ , R ₂ N ⁻)	Not Important: Usually Weak (ROH, R ₂ NH)
Leaving Group:	Moderately Important (same trend as S _N 1)	Very Important (same trend as S _N 1)
Solvent:	Wide Range of Solvents	Polar Protic
Product Ratio:	<p>Zaitsev Rule: The most highly substituted alkene usually predominates.</p> <p>Hofmann Product: Use of a sterically hindered base will result in formation of the least substituted alkene (Hofmann product).</p>	

Generic Reaction-Energy Diagrams

Substitution





Elimination




Bimolecular

Unimolecular



Predicting the Products: Substitution versus Elimination



* Under conditions that favor a unimolecular reaction (weak nuc/base and polar protic solvent), mixtures of S_N1 and $E1$ are usually obtained.