

Lecture notes in Fluid Dynamics
(1.63J/2.01J)
by Chiang C. Mei, MIT

4-3MTwind.tex

4.3 Buoyancy-driven convection - The Valley Wind

ref: Prandtl: **Fluid Dynamics**.

Due to solar heating during the day, a mountain slope may be warmer than the surrounding air in a summer night. Let the air near a mountain slope be stably stratified

$$T_o = T_0 + Ny', \quad (4.3.1)$$

where $T_0 = \text{constant}$, and $N > 0$. Let the slope temperature be :

$$T_s = T_1 + Ny', \quad (4.3.2)$$

where $T_1 > T_0$. See the left of Figure 4.3.2. Consider first the static equilibrium:




Figure 4.3.1: Thermal convection along a slope

$$0 = -\frac{dp_o}{dz} - \rho_o g$$

hence

$$p_o = p_o(\infty) + \int_z^{\infty} \rho_o g dz$$

Let A and B be two points with the same elevation but A is on the slope and B is in the air. Since $p_A < p_B$,

$$\frac{\partial p_o}{\partial x} < 0$$

implying

$$\frac{\partial p_o}{\partial x'} < 0$$

The pressure gradient must drive an upward flow along the slope.

Let us consider the dynamics. Let

$$T(x, y) = T_o + \theta(y) \quad (4.3.3)$$

and

$$\rho(x, y) = \rho_o + S(y) = \text{static density} + \text{dynamic density} \quad (4.3.4)$$

By the equation of state,

$$\rho = \rho_o [1 - \beta (T - T_o)] = \rho_o [1 - \beta (T_o - T_o)] - \rho_o \beta \theta.$$

Therefore

$$\rho_o = \rho_o [1 - \beta (T_o - T_o)] = \rho_o (1 - \beta N y') \quad (4.3.5)$$

and

$$S(x, y) = -\rho_o \beta \theta(x, y) \quad (4.3.6)$$

Note by rotation of coordinates,

$$T_o - T_0 = N y' = N(x \sin \alpha + y \cos \alpha). \quad (4.3.7)$$

The flow equations are:

$$u_x + v_y = 0 \quad (4.3.8)$$

$$\rho (u u_x + v u_y) = -p_{dx} + \mu (u_{xx} + u_{yy}) - (\rho - \rho_a) g \sin \alpha \quad (4.3.9)$$

$$\rho (u v_x + v v_u) = -p_{dy} + \mu (v_{xx} + v_{yy}) - (\rho - \rho_a) g \cos \alpha \quad (4.3.10)$$

$$u T_x + v T_y = k (T_{xx} + T_{yy}), \quad (4.3.11)$$

where T is the total temperature and

$$k = \frac{K}{\bar{\rho}_o c_p}.$$

is the thermal diffusivity. Since $\partial/\partial x = 0$, $v = 0$ from continuity. From Eqn. (4.3.9)

$$\nu u_{yy} + (\beta g \sin \alpha) \theta = 0. \quad (4.3.12)$$

after invoking Boussinesq approximation. In Eqn. (4.3.11),

$$\frac{\partial T}{\partial x} = \frac{\partial T_o}{\partial x} = N \sin \alpha.$$

Therefore,

$$u N \sin \alpha = k \theta_{yy}. \quad (4.3.13)$$

Combining Eqns. (4.3.12) and (4.3.13), we get

$$\frac{d^4 u}{dy^4} + \left(\frac{\beta g N \sin^2 \alpha}{\nu k} \right) u = 0 \quad (4.3.14)$$

and

$$\frac{d^4 \theta}{dy^4} + \left(\frac{\beta g N \sin^2 \alpha}{\nu k} \right) \theta = 0 \quad (4.3.15)$$

Let

$$\ell^4 = \frac{4\nu k}{\beta g A \sin^2 \alpha} \quad \text{and} \quad y = \ell \eta \quad (4.3.16)$$

then

$$\frac{d^4 u}{d\eta^4} + 4u = 0; \quad \text{and} \quad \frac{d^4 \theta}{d\eta^4} + 4\theta = 0 \quad (4.3.17)$$

The velocity is

$$u = U e^{-\eta} \sin \eta \quad \text{so that} \quad u(0) = 0 \quad (4.3.18)$$

The temperature is

$$\theta = \theta_0 e^{-\eta} \cos \eta \quad (4.3.19)$$

The boundary conditions at $\eta \sim \infty$ are satisfied. In order that $\theta(0) = T_1 - T_0$ on $\eta = 0$ we choose

$$\theta_0 = T_1 - T_0 \quad (4.3.20)$$

Note that the boundary layer thickness is

$$\delta \sim O(\ell) \sim \left(\frac{4\nu k}{\rho g N \sin^2 \alpha} \right)^{1/4} \quad (4.3.21)$$

Thus if $\alpha \downarrow, \delta \uparrow$ as $1/\sin^2 \alpha$.

Using Eqn. (4.3.13), we get

$$N \sin \alpha U e^{-\eta} \sin \eta = k \left(\frac{\beta g N \sin^2 \alpha}{4\nu k} \right)^{1/2} 2\theta_0 e^{-\eta} \sin \eta.$$

Hence,

$$U = \theta_0 \left(\frac{\beta g k}{N \nu} \right)^{1/2} \quad (4.3.22)$$

Finally

$$u = (T_1 - T_0) \left(\frac{\beta g k}{N \nu} \right)^{1/2} e^{-\eta} \sin \eta. \quad (4.3.23)$$

and

$$\theta = (T_1 - T_0) e^{-\eta} \cos \eta \quad (4.3.24)$$

It is easy to show from (4.3.13) that the total mass flux rate is

$$M = \int_0^\infty \rho u dy = \rho_0 \beta k \left. \frac{d\theta}{dy} \right|_0. \quad (4.3.25)$$

Note from (4.3.22) that U is independent of α . If $\alpha \downarrow$, the buoyancy force is weaker, but the shear rate $\partial u/\partial y$ is smaller, hence the wall resistance is smaller. U is not reduced!




Figure 4.3.2: Wind along a valley due to feeding from mountains

On a warm slope (due to solar heating during the day) , air rises at night. If there are two slopes forming a valley, fluid must be supplied from the bottom of the valley; this is the reason for valley wind blowing from low altitude to high.

On a cold slope (due to radiation loss at night) air sinks at high noon. Valley wind must flow from high to low.