

Quiz 2 Solutions:
November 3, 2009

Problem 2. (49 points)

(a) (7 points)

We start by recognizing that $f_X(x) = e^{-x}$ for $x \geq 0$ and $f_{Y|X}(y | x) = xe^{-xy}$ for $y \geq 0$. Furthermore, $f_{X,Y}(x, y) = f_X(x) \cdot f_{Y|X}(y | x)$. Substituting for $f_X(x)$ and $f_{Y|X}(y | x)$ yields,

$$f_{X,Y}(x, y) = \begin{cases} xe^{-(1+y)x}, & x \geq 0, y \geq 0 \\ 0, & \text{otherwise.} \end{cases}$$

(b) (7 points)

The marginal PDF of Y can be found by integrating the joint PDF of X and Y .

$$\begin{aligned} f_Y(y) &= \int_X f_{X,Y}(x, y) dx \\ &= \int_0^\infty xe^{-(1+y)x} dx \end{aligned}$$

$$f_Y(y) = \begin{cases} \frac{1}{(1+y)^2}, & y \geq 0 \\ 0, & \text{otherwise.} \end{cases}$$

(c) (7 points)

We are asked to compute the PDF of the random variable X while conditioning on another random variable Y . The conditional PDF of X given that $Y = 2$ is

$$f_{X|Y}(x | 2) = \frac{f_{X,Y}(x, 2)}{f_Y(2)} = \frac{xe^{-3x}}{\frac{1}{3^2}}$$

$$f_{X|Y}(x | 2) = \begin{cases} 9xe^{-3x}, & x \geq 0 \\ 0, & \text{otherwise.} \end{cases}$$

(d) (7 points)

$$\begin{aligned} \mathbf{E}[X | Y = 2] &= \int_X x \cdot f_{X|Y}(x | 2) dx \\ &= 9 \int_0^\infty x^2 e^{-3x} dx \\ &= 9 \cdot \frac{2}{3^3} \\ &= \frac{2}{3}. \end{aligned}$$

(e) (7 points)

In the new universe in which $X = 2$, we are asked to compute the conditional PDF of Y given the event $Y \geq 3$.

$$f_{Y|X, Y \geq 3}(y | 2) = \frac{f_{Y|X}(y | 2)}{\mathbf{P}(Y \geq 3 | X = 2)}.$$

We first calculate the $\mathbf{P}(Y \geq 3 | X = 2)$.

$$\begin{aligned} \mathbf{P}(Y \geq 3 | X = 2) &= \int_3^{\infty} f_{Y|X}(y | 2) dy \\ &= \int_3^{\infty} 2e^{-2y} dy \\ &= 1 - F_{Y|X}(3 | 2) \\ &= 1 - (1 - e^{-2 \cdot 3}) \\ &= e^{-6}, \end{aligned}$$

where $F_{Y|X}(3|2)$ is the CDF of an exponential random variable with $\lambda = 2$ evaluated at $y = 3$. Substituting the values of $f_{Y|X}(y | 2)$ and $\mathbf{P}(Y \geq 3 | X = 2)$ yields

$$f_{Y|X, Y \geq 3}(y | 2) = \begin{cases} 2e^6 e^{-2y}, & y \geq 3 \\ 0, & \text{otherwise.} \end{cases}$$

Alternatively, $f_{Y|X}(y | 2)$ is an exponential random variable with $\lambda = 2$. To compute the conditional PMF $f_{Y|X, Y \geq 3}(y | 2)$, we can apply the memorylessness property of an exponential variable. Therefore, this conditional PMF is also an exponential random variable with $\lambda = 2$, but it is shifted by 3.

(f) (7 points)

Let's define $Z = e^{2X}$. Since X is an exponential random variable that takes on non-negative values ($X \geq 0$), $Z \geq 1$. We find the PDF of Z by first computing its CDF.

$$\begin{aligned} F_Z(z) &= \mathbf{P}(Z \leq z) \\ &= \mathbf{P}(e^{2X} \leq z) \\ &= \mathbf{P}(2X \leq \ln z) \\ &= \mathbf{P}\left(X \leq \frac{\ln z}{2}\right) \\ &= 1 - e^{-\frac{\ln z}{2}} \\ &= 1 - e^{\ln z^{-\frac{1}{2}}} \end{aligned}$$

The CDF of Z is:

$$F_Z(z) = \begin{cases} 1 - z^{-\frac{1}{2}} & z \geq 1 \\ 0, & z < 1 \end{cases}$$

Differentiating the CDF of Z yields the PDF

$$f_Z(z) = \begin{cases} \frac{1}{2}z^{-\frac{3}{2}} & z \geq 1 \\ 0, & z < 1 \end{cases}$$

Alternatively, you can apply the PDF formula for a strictly monotonic function of a continuous random variable. Recall if $z = g(x)$ and $x = h(z)$, then

$$f_Z(z) = f_X(h(z)) \left| \frac{dh}{dz}(z) \right|.$$

In this problem, $z = e^{2x}$ and $x = \frac{1}{2}\ln z$. Note that $f_Z(z)$ is nonzero for $z > 1$. Since X is an exponential random variable with $\lambda = 1$, $f_X(x) = e^{-x}$. Thus,

$$\begin{aligned} f_Z(z) &= e^{-\frac{1}{2}\ln z} \left| \frac{1}{2z} \right| \\ &= e^{\ln z^{-\frac{1}{2}}} \frac{1}{2z} \\ &= \frac{1}{2}z^{-\frac{3}{2}} \quad z \geq 1, \end{aligned}$$

where the second equality holds since the expression inside the absolute value is always positive for $z \geq 1$.

Problem 3. (10 points)

(a) (5 points) The quantity $\mathbf{E}[X | Y]$ is always:

- (i) A number.
- (ii) A discrete random variable.
- (iii) A continuous random variable.
- (iv) Not enough information to choose between (i)-(iii).

If X and Y are not independent, then $\mathbf{E}[X | Y]$ is a function of Y and is therefore a continuous random variable. However if X and Y are independent, then $\mathbf{E}[X | Y] = \mathbf{E}[X]$ which is a number.

(b) (5 points) The quantity $\mathbf{E}[\mathbf{E}[X | Y, N] | N]$ is always:

- (i) A number.
- (ii) A discrete random variable.
- (iii) A continuous random variable.
- (iv) Not enough information to choose between (i)-(iii).

If X , Y and N are not independent, then the inner expectation $G(Y, N) = \mathbf{E}[X | Y, N]$ is a function of Y and N . Furthermore $\mathbf{E}[G(Y, N) | N]$ is a function of N , a discrete random variable. If X , Y and N are independent, then the inner expectation $\mathbf{E}[X | Y, N] = \mathbf{E}[X]$, which is a number. The expectation of a number given N is still a number, which is a special case of a discrete random variable.

Problem 4. (25 points)

(a) (i) (5 points)

Using the Law of Iterated Expectations, we have

$$\mathbf{E}[X] = \mathbf{E}[\mathbf{E}[X | Q]] = \mathbf{E}[Q] = \frac{1}{2}.$$

(ii) (5 points)

X is a Bernoulli random variable with a mean $p = \frac{1}{2}$ and its variance is $\text{var}(X) = p(1 - p) = 1/4$.

(b) (7 points)

We know that $\text{cov}(X, Q) = \mathbf{E}[XQ] - \mathbf{E}[X]\mathbf{E}[Q]$, so first let's calculate $\mathbf{E}[XQ]$:

$$\mathbf{E}[XQ] = \mathbf{E}[\mathbf{E}[XQ | Q]] = \mathbf{E}[Q\mathbf{E}[X | Q]] = \mathbf{E}[Q^2] = \frac{1}{3}.$$

Therefore, we have

$$\text{cov}(X, Q) = \frac{1}{3} - \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{12}.$$

(c) (8 points)

Using Bayes' Rule, we have

$$f_{Q|X}(q | 1) = \frac{f_Q(q)p_{X|Q}(1 | q)}{p_X(1)} = \frac{f_Q(q)\mathbf{P}(X = 1 | Q = q)}{\mathbf{P}(X = 1)}, \quad 0 \leq q \leq 1.$$

Additionally, we know that

$$\mathbf{P}(X = 1 | Q = q) = q,$$

and that for Bernoulli random variables

$$\mathbf{P}(X = 1) = \mathbf{E}[X] = \frac{1}{2}.$$

Thus, the conditional PDF of Q given $X = 1$ is

$$\begin{aligned} f_{Q|X}(q | 1) &= \frac{1 \cdot q}{1/2} \\ &= \begin{cases} 2q, & 0 \leq q \leq 1, \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

Problem 5. (21 points)

(a) (7 points)

$$\begin{aligned} \mathbf{P}(S \geq 1) &= \mathbf{P}(\min\{X, Y\} \geq 1) = \mathbf{P}(X \geq 1 \text{ and } Y \geq 1) = \mathbf{P}(X \geq 1)\mathbf{P}(Y \geq 1) \\ &= (1 - F_X(1))(1 - F_Y(1)) = (1 - \Phi(1))^2 \approx (1 - 0.8413)^2 \approx 0.0252. \end{aligned}$$

(b) (7 points)

Recalling Problem 2 of Problem Set 6, we have

$$\begin{aligned}\mathbf{P}(s \leq S \text{ and } L \leq \ell) &= \mathbf{P}(s \leq \min\{X, Y\} \text{ and } \max\{X, Y\} \leq \ell) \\ &= \mathbf{P}(s \leq X \text{ and } s \leq Y \text{ and } X \leq \ell \text{ and } Y \leq \ell) \\ &= \mathbf{P}(s \leq X \leq \ell)\mathbf{P}(s \leq Y \leq \ell) \\ &= (F_X(\ell) - F_X(s))(F_Y(\ell) - F_Y(s)).\end{aligned}$$

(c) (7 points)

Given that $s \leq s + \delta \leq \ell$, the event $\{s \leq S \leq s + \delta, \ell \leq L \leq \ell + \delta\}$ is made up of the union of two disjoint possible events:

$$\{s \leq X \leq s + \delta, \ell \leq Y \leq \ell + \delta\} \cup \{s \leq Y \leq s + \delta, \ell \leq X \leq \ell + \delta\}.$$

In other words, either $S = X$ and $L = Y$, or $S = Y$ and $L = X$. Because the two events are disjoint, the probability of their union is equal to the sum of their individual probabilities.

Using also the independence of X and Y , we have

$$\begin{aligned}\mathbf{P}(s \leq S \leq s + \delta, \ell \leq L \leq \ell + \delta) &= \mathbf{P}(s \leq X \leq s + \delta, \ell \leq Y \leq \ell + \delta) \\ &\quad + \mathbf{P}(s \leq Y \leq s + \delta, \ell \leq X \leq \ell + \delta) \\ &= \mathbf{P}(s \leq X \leq s + \delta)\mathbf{P}(\ell \leq Y \leq \ell + \delta) \\ &\quad + \mathbf{P}(s \leq Y \leq s + \delta)\mathbf{P}(\ell \leq X \leq \ell + \delta) \\ &= \int_s^{s+\delta} f_X(x)dx \int_\ell^{\ell+\delta} f_Y(y)dy \\ &\quad + \int_s^{s+\delta} f_Y(y)dy \int_\ell^{\ell+\delta} f_X(x)dx\end{aligned}$$

MIT OpenCourseWare
<http://ocw.mit.edu>

6.041 / 6.431 Probabilistic Systems Analysis and Applied Probability
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.