

**Tutorial 5: Solutions**

1. (a) Let  $A$  be the event that the machine is functional. Conditioned on the random variable  $Q$  taking on a particular value  $q$ ,  $\mathbf{P}(A|Q = q) = q$ . Using the continuous form of the total probability theorem, the probability of event  $A$  is given by:

$$\begin{aligned}\mathbf{P}(A) &= \int_0^1 \mathbf{P}(A|Q = q)f_Q(q)dq \\ &= \int_0^1 q dq \\ &= 1/2\end{aligned}$$

- (b) Let  $B$  be the event that the machine is functional on  $m$  out of the last  $n$  days. Conditioned on random variable  $Q$  taking on value  $q$  (a probability  $q$  of being functional) the probability of event  $B$  is binomial with  $n$  trials,  $m$  successes, and a probability  $q$  of success in each trial. Again using the total probability theorem, the probability of event  $B$  is given by:

$$\begin{aligned}\mathbf{P}(B) &= \int_0^1 \mathbf{P}(B|Q = q)f_Q(q)dq \\ &= \int_0^1 \binom{n}{m} q^m (1 - q)^{n-m} f_Q(q) dq \\ &= \binom{n}{m} \frac{m!(n-m)!}{(n+1)!}\end{aligned}$$

We then find the distribution on  $Q$  conditioned on event  $B$  using Bayes rule:

$$\begin{aligned}f_{Q|B}(q) &= \frac{\mathbf{P}(B|Q = q)f_Q(q)}{\mathbf{P}(B)} \\ &= \frac{q^m (1 - q)^{n-m}}{\frac{m!(n-m)!}{(n+1)!}} \quad 0 \leq q \leq 1, \quad n \geq m.\end{aligned}$$

2. Since  $Y = |X|$  you can visualize the PDF for any given  $y$  as

$$f_Y(y) = \begin{cases} f_X(y) + f_X(-y), & \text{if } y \geq 0, \\ 0, & \text{if } y < 0. \end{cases}$$

Also note that since  $Y = |X|$ ,  $Y \geq 0$ .

(a)  $f_X(x) = \begin{cases} \frac{1}{3}, & \text{if } -2 < x \leq 1, \\ 0, & \text{otherwise.} \end{cases}$

So,  $f_X(x)$  for  $-1 \leq x \leq 0$  gets added to  $f_X(x)$  for  $0 \leq x \leq 1$ :

$$f_Y(y) = \begin{cases} 2/3, & \text{if } 0 \leq y \leq 1, \\ 1/3, & \text{if } 1 < y \leq 2, \\ 0, & \text{otherwise.} \end{cases}$$

MASSACHUSETTS INSTITUTE OF TECHNOLOGY  
 Department of Electrical Engineering & Computer Science  
**6.041/6.431: Probabilistic Systems Analysis**  
 (Fall 2010)

---

(b) Here we are told  $X > 0$ . So there are no negative values of  $X$  that need to be considered. Thus,

$$f_Y(y) = f_X(y) = \begin{cases} 2e^{-2y}, & \text{if } y \geq 0, \\ 0, & \text{otherwise.} \end{cases}$$


(c) As explained in the beginning,  $f_Y(y) = f_X(y) + f_X(-y)$ .

3. We want to compute the CDF of the ambulance's travel time  $T$ ,  $\mathbf{P}(T \leq t) = \mathbf{P}(|X - Y| \leq vt)$ , where  $X$  and  $Y$  are the locations of the ambulance and accident (uniform over  $[0, l]$ ). Since  $X$  and  $Y$  are independent, we know:

$$f_{X,Y}(x, y) = \begin{cases} \frac{1}{l^2} & , \text{ if } 0 \leq x, y \leq l \\ 0 & , \text{ otherwise} \end{cases} .$$

$$\begin{aligned} \mathbf{P}(T \leq t) &= \mathbf{P}(|X - Y| \leq vt) = \mathbf{P}(-vt \leq Y - X \leq vt) \\ &= \mathbf{P}(X - vt \leq Y \leq X + vt) \end{aligned}$$

We can see that  $\mathbf{P}(X - vt \leq Y \leq X + vt)$  corresponds to the integral of the joint density of  $X$  and  $Y$  over the shaded region in the figure below:



Therefore, because the joint density is uniform over the entire region, we have:

$$F_T(t) = (1/l^2) \times (\text{Shaded area}) = \begin{cases} 0 & , \text{ if } t < 0 \\ \frac{2vt}{l} - \frac{(vt)^2}{l^2} & , \text{ if } 0 \leq t < \frac{l}{v} \\ 1 & , \text{ if } t \geq \frac{l}{v} \end{cases} .$$

By differentiating the CDF, we find the density of  $T$ :

$$f_T(t) = \begin{cases} \frac{2v}{l} - \frac{2v^2t}{l^2} & , \text{ if } 0 \leq t \leq \frac{l}{v} \\ 0 & , \text{ otherwise} \end{cases} .$$

MIT OpenCourseWare  
<http://ocw.mit.edu>

6.041 / 6.431 Probabilistic Systems Analysis and Applied Probability  
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.