

**6.041 Probabilistic Systems Analysis
6.431 Applied Probability**

- Staff:
 - Lecturer: John Tsitsiklis
- Pick up **and read** course information handout
- **Turn in recitation and tutorial scheduling form** (last sheet of course information handout)
- Pick up copy of slides

Coursework

- Quiz 1 (October 12, 12:05-12:55pm) 17%
- Quiz 2 (November 2, 7:30-9:30pm) 30%
- Final exam (scheduled by registrar) 40%
- Weekly homework (best 9 of 10) 10%
- Attendance/participation/enthusiasm in recitations/tutorials 3%
- **Collaboration policy** described in course info handout
- Text: *Introduction to Probability*, 2nd Edition, D. P. Bertsekas and J. N. Tsitsiklis, Athena Scientific, 2008
Read the text!

LECTURE 1

- **Readings:** Sections 1.1, 1.2

Lecture outline


- Probability as a mathematical framework for reasoning about uncertainty
- Probabilistic models
 - sample space
 - probability law
- Axioms of probability
- Simple examples

Sample space Ω

- “List” (set) of possible outcomes
- List must be:
 - Mutually exclusive
 - Collectively exhaustive
- Art: to be at the “right” granularity


Sample space: Discrete example

- Two rolls of a tetrahedral die
- Sample space vs. sequential description



Sample space: Continuous example

$$\Omega = \{(x, y) \mid 0 \leq x, y \leq 1\}$$



Probability axioms

- **Event:** a subset of the sample space
- Probability is assigned to events


Axioms:

1. **Nonnegativity:** $P(A) \geq 0$
2. **Normalization:** $P(\Omega) = 1$
3. **Additivity:** If $A \cap B = \emptyset$, then $P(A \cup B) = P(A) + P(B)$

$$P(\{s_1, s_2, \dots, s_k\}) = P(\{s_1\}) + \dots + P(\{s_k\}) \\ = P(s_1) + \dots + P(s_k)$$

- Axiom 3 needs strengthening
- Do weird sets have probabilities?

Probability law: Example with finite sample space



- Let every possible outcome have probability 1/16
- $P((X, Y) \text{ is } (1,1) \text{ or } (1,2)) =$
- $P(\{X = 1\}) =$
- $P(X + Y \text{ is odd}) =$
- $P(\min(X, Y) = 2) =$

Discrete uniform law


- Let all outcomes be equally likely
- Then,

$$P(A) = \frac{\text{number of elements of } A}{\text{total number of sample points}}$$

- Computing probabilities \equiv counting
- Defines fair coins, fair dice, well-shuffled decks

Continuous uniform law


- Two “random” numbers in $[0, 1]$.



- **Uniform** law: Probability = Area
 - $P(X + Y \leq 1/2) = ?$
 - $P((X, Y) = (0.5, 0.3))$

Probability law: Ex. w/countably infinite sample space

- Sample space: $\{1, 2, \dots\}$
 - We are given $P(n) = 2^{-n}$, $n = 1, 2, \dots$
 - Find $P(\text{outcome is even})$



$$P(\{2, 4, 6, \dots\}) = P(2) + P(4) + \dots = \frac{1}{2^2} + \frac{1}{2^4} + \frac{1}{2^6} + \dots = \frac{1}{3}$$

- **Countable additivity axiom** (needed for this calculation):
If A_1, A_2, \dots are disjoint events, then:

$$P(A_1 \cup A_2 \cup \dots) = P(A_1) + P(A_2) + \dots$$

Remember!

- Turn in recitation/tutorial scheduling form **now**
- Tutorials start next week

MIT OpenCourseWare
<http://ocw.mit.edu>

6.041 / 6.431 Probabilistic Systems Analysis and Applied Probability
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.