

## LECTURE 17

### Markov Processes – II

- **Readings:** Section 7.3


#### Lecture outline

- Review
- Steady-State behavior
  - Steady-state convergence theorem
  - Balance equations
- Birth-death processes

## Review

- Discrete state, discrete time, time-homogeneous
  - Transition probabilities  $p_{ij}$
  - Markov property
- $r_{ij}(n) = \mathbf{P}(X_n = j \mid X_0 = i)$
- Key recursion:
 
$$r_{ij}(n) = \sum_k r_{ik}(n-1)p_{kj}$$

### Warmup



$$\mathbf{P}(X_1 = 2, X_2 = 6, X_3 = 7 \mid X_0 = 1) =$$



$$\mathbf{P}(X_4 = 7 \mid X_0 = 2) =$$

### Recurrent and transient states

- State  $i$  is **recurrent** if:
  - starting from  $i$ ,
  - and from wherever you can go,
  - there is a way of returning to  $i$
- If not recurrent, called **transient**
- **Recurrent class:**
  - collection of recurrent states that
  - “communicate” to each other
  - and to no other state

### Periodic states

- The states in a recurrent class are **periodic** if they can be grouped into  $d > 1$  groups so that all transitions from one group lead to the next group



### Steady-State Probabilities

- Do the  $r_{ij}(n)$  converge to some  $\pi_j$ ? (independent of the initial state  $i$ )
- Yes, if:
  - recurrent states are all in a single class, and
  - single recurrent class is not periodic
- Assuming “yes,” start from key recursion

$$r_{ij}(n) = \sum_k r_{ik}(n-1)p_{kj}$$

- take the limit as  $n \rightarrow \infty$

$$\pi_j = \sum_k \pi_k p_{kj}, \quad \text{for all } j$$


- Additional equation:

$$\sum_j \pi_j = 1$$


### Visit frequency interpretation

$$\pi_j = \sum_k \pi_k p_{kj}$$


- (Long run) frequency of being in  $j$ :  $\pi_j$
- Frequency of transitions  $k \rightarrow j$ :  $\pi_k p_{kj}$
- Frequency of transitions into  $j$ :  $\sum_k \pi_k p_{kj}$



### Example



### Birth-death processes



- Special case:  $p_i = p$  and  $q_i = q$  for all  $i$   
 $\rho = p/q = \text{load factor}$

$$\pi_{i+1} = \pi_i \frac{p}{q} = \pi_i \rho$$

$$\pi_i = \pi_0 \rho^i, \quad i = 0, 1, \dots, m$$

- Assume  $p < q$  and  $m \approx \infty$

$$\pi_0 = 1 - \rho$$

$$\mathbf{E}[X_n] = \frac{\rho}{1 - \rho} \quad (\text{in steady-state})$$

MIT OpenCourseWare  
<http://ocw.mit.edu>

6.041 / 6.431 Probabilistic Systems Analysis and Applied Probability  
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.