

LECTURE 3

- **Readings:** Section 1.5
- Review
- Independence of two events
- Independence of a collection of events

Review

$$P(A | B) = \frac{P(A \cap B)}{P(B)}, \quad \text{assuming } P(B) > 0$$

- Multiplication rule:

$$P(A \cap B) = P(B) \cdot P(A | B) = P(A) \cdot P(B | A)$$

- Total probability theorem:

$$P(B) = P(A)P(B | A) + P(A^c)P(B | A^c)$$


- Bayes rule:

$$P(A_i | B) = \frac{P(A_i)P(B | A_i)}{P(B)}$$

Models based on conditional probabilities

- 3 tosses of a biased coin:

$$P(H) = p, \quad P(T) = 1 - p$$



$$P(THT) =$$

$$P(1 \text{ head}) =$$

$$P(\text{first toss is H} | 1 \text{ head}) =$$


Independence of two events

- **“Defn:”** $P(B | A) = P(B)$
 - “occurrence of A provides no information about B 's occurrence”
- Recall that $P(A \cap B) = P(A) \cdot P(B | A)$
- **Defn:** $P(A \cap B) = P(A) \cdot P(B)$
- Symmetric with respect to A and B
 - applies even if $P(A) = 0$
 - implies $P(A | B) = P(A)$

Conditioning may affect independence

- Conditional independence, given C , is defined as independence under probability law $P(\cdot | C)$


- Assume A and B are independent



- If we are told that C occurred, are A and B independent?

Conditioning may affect independence

- Two unfair coins, A and B :
 $P(H \mid \text{coin } A) = 0.9$, $P(H \mid \text{coin } B) = 0.1$
 choose either coin with equal probability



- Once we know it is coin A , are tosses independent?
- If we do not know which coin it is, are tosses independent?
 - Compare:
 - $P(\text{toss } 11 = H)$
 - $P(\text{toss } 11 = H \mid \text{first } 10 \text{ tosses are heads})$

Independence of a collection of events

- Intuitive definition:
 Information on some of the events tells us nothing about probabilities related to the remaining events
 - E.g.:
 $P(A_1 \cap (A_2^c \cup A_3) \mid A_5 \cap A_6^c) = P(A_1 \cap (A_2^c \cup A_3))$
- Mathematical definition:
 Events A_1, A_2, \dots, A_n are called **independent** if:

$$P(A_i \cap A_j \cap \dots \cap A_q) = P(A_i)P(A_j) \dots P(A_q)$$
 for any distinct indices i, j, \dots, q ,
 (chosen from $\{1, \dots, n\}$)

Independence vs. pairwise independence

- Two independent fair coin tosses
 - A : First toss is H
 - B : Second toss is H
 - $P(A) = P(B) = 1/2$

HH	HT
TH	TT

- C : First and second toss give same result
- $P(C) =$
- $P(C \cap A) =$
- $P(A \cap B \cap C) =$
- $P(C \mid A \cap B) =$
- Pairwise independence **does not** imply independence

The king's sibling

- The king comes from a family of two children. What is the probability that his sibling is female?

MIT OpenCourseWare
<http://ocw.mit.edu>

6.041 / 6.431 Probabilistic Systems Analysis and Applied Probability
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.