

## LECTURE 4

- **Readings:** Section 1.6

### Lecture outline

- Principles of counting
  - permutations
  - $k$ -permutations
  - combinations
  - partitions
- Binomial probabilities

## Discrete uniform law


- Let all sample points be equally likely
- Then,

$$P(A) = \frac{\text{number of elements of } A}{\text{total number of sample points}} = \frac{|A|}{|\Omega|}$$

- Just count...

## Basic counting principle

- $r$  stages
- $n_i$  choices at stage  $i$



- **Number of choices is:**  $n_1 n_2 \cdots n_r$
- Number of license plates with 3 letters and 4 digits =
- ... if repetition is prohibited =
- **Permutations:** Number of ways of ordering  $n$  elements is:
- Number of subsets of  $\{1, \dots, n\}$  =

## Example

- Probability that six rolls of a six-sided die all give different numbers?
  - Number of outcomes that make the event happen:
  - Number of elements in the sample space:
  - Answer:

### Combinations

- $\binom{n}{k}$ : number of  $k$ -element subsets of a given  $n$ -element set
- Two ways of constructing an ordered sequence of  $k$  **distinct** items:
  - Choose the  $k$  items one at a time:  
 $n(n-1)\cdots(n-k+1) = \frac{n!}{(n-k)!}$  choices
  - Choose  $k$  items, then order them ( $k!$  possible orders)

• Hence:

$$\binom{n}{k} \cdot k! = \frac{n!}{(n-k)!}$$

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

$$\sum_{k=0}^n \binom{n}{k} =$$

### Binomial probabilities

- $n$  independent coin tosses
  - $P(H) = p$
- $P(HTTTHHH) =$
- $P(\text{sequence}) = p^{\# \text{ heads}}(1-p)^{\# \text{ tails}}$

$$\begin{aligned} P(k \text{ heads}) &= \sum_{k\text{-head seq.}} P(\text{seq.}) \\ &= (\# \text{ of } k\text{-head seqs.}) \cdot p^k(1-p)^{n-k} \\ &= \binom{n}{k} p^k(1-p)^{n-k} \end{aligned}$$

### Coin tossing problem

- event  $B$ : 3 out of 10 tosses were “heads”.
  - Given that  $B$  occurred, what is the (conditional) probability that the first 2 tosses were heads?
- All outcomes in set  $B$  are equally likely: probability  $p^3(1-p)^7$ 
  - Conditional probability law is uniform
- Number of outcomes in  $B$ :
- Out of the outcomes in  $B$ , how many start with HH?

### Partitions

- 52-card deck, dealt to 4 players
- Find  $P(\text{each gets an ace})$
- Outcome: a partition of the 52 cards
  - number of outcomes:
- Count number of ways of distributing the four aces:  $4 \cdot 3 \cdot 2$
- Count number of ways of dealing the remaining 48 cards
- Answer:

$$\frac{52!}{13! 13! 13! 13!}$$

$$\frac{48!}{12! 12! 12! 12!}$$

$$\frac{4 \cdot 3 \cdot 2 \cdot \frac{48!}{12! 12! 12! 12!}}{\frac{52!}{13! 13! 13! 13!}}$$

MIT OpenCourseWare  
<http://ocw.mit.edu>

6.041SC Probabilistic Systems Analysis and Applied Probability  
Fall 2013

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.