

**Recitation 4 Solutions**  
**September 21, 2010**

1. The sample space consists of all possible choices for the birthday of each person. Since there are  $n$  persons, and each has 365 choices for their birthday, the sample space has  $365^n$  elements. Let us now consider those choices of birthdays for which no two persons have the same birthday. Assuming that  $n \leq 365$ , there are 365 choices for the first person, 364 for the second, etc., for a total of  $365 \cdot 364 \cdots (365 - n + 1)$ . Thus,

$$\mathbf{P}(\text{no two birthdays coincide}) = \frac{365 \cdot 364 \cdots (365 - n + 1)}{365^n}.$$

It is interesting to note that for  $n$  as small as 23, the probability that there are two persons with the same birthday is larger than  $1/2$ .




Image by MIT OpenCourseWare.

2. As we have done before, we will count the number of favorable positions in which we can safely place 8 rooks, and then divide this by the total number of positions for 8 rooks on a  $8 \times 8$  chessboard. First we count the number of favorable positions for the rooks. We will place the

MASSACHUSETTS INSTITUTE OF TECHNOLOGY  
Department of Electrical Engineering & Computer Science  
**6.041/6.431: Probabilistic Systems Analysis**  
(Fall 2010)

---

rooks one by one. For the first rook, there are no constraints, so we have 64 choices. Placing this rook, however, eliminates one row and one column. Thus for our second rook, we can imagine that the illegal column and row have been removed, thus leaving us with a  $7 \times 7$  chessboard, and thus with 49 choices. Similarly, for the third rook we have 36 choices, for the fourth 25, etc... There are  $64 \cdot 49 \cdot 36 \cdot 25 \cdot 16 \cdot 9 \cdot 4$  total ways we can place 8 rooks without any restrictions, and therefore the probability we are after is:

$$\frac{64 \cdot 49 \cdot 36 \cdot 25 \cdot 16 \cdot 9 \cdot 4}{\frac{64!}{56!}}$$

- See textbook, Problem 1.61, page 69.
- The group of  $n$  slots is divided into segments of length  $n_1, \dots, n_r$  slots. The  $n$  items can be arranged in  $n!$  ways, where each arrangement corresponds to a partition into the  $r$  segments. But all arrangements within a single segment lead to the same partition, where there are  $n_i!$  ways to arrange the items within  $i$ th segment. Thus, for each segment we must divide by the number of ways to arrange the items within that segment. The solution is then:

$$\frac{\text{Ways to arrange } n \text{ items}}{(\text{Ways to arrange items in segment 1}) \cdots (\text{Ways to arrange items in segment } r)} = \frac{n!}{n_1! \cdots n_r!}$$

- The probability of drawing a particular sequence of balls containing exactly  $n_i$  of color  $i$  balls is  $p_1^{n_1} \cdots p_r^{n_r}$ . The number of possible sequences containing  $n_i$  of color  $i$  balls is the number of ways to form a partition of  $n$  distinct slots into subsets of cardinality  $n_1, \dots, n_r$  which is  $\binom{n}{n_1, \dots, n_r}$ . Therefore, the probability of obtaining exactly  $n_i$  balls of color  $i$  is:

$$\binom{n}{n_1, \dots, n_r} p_1^{n_1} \cdots p_r^{n_r}$$

MIT OpenCourseWare  
<http://ocw.mit.edu>

6.041SC Probabilistic Systems Analysis and Applied Probability  
Fall 2013

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.