

## In-Class Problems Week 11, Fri.

**Problem 1.** (a) Verify that

$$\frac{1}{(1-x)^k} = \sum_{n=0}^{\infty} \binom{n+k-1}{n} x^n.$$

*Hint:* Use the fact that if  $A(x) = \sum_{n=0}^{\infty} a_n x^n$ , then

$$a_n = \frac{A^{(n)}(0)}{n!},$$

where  $A^{(n)}$  is the  $n$ th derivative of  $A$ .

(b) Let  $S(x) ::= \sum_{k=1}^{\infty} k^2 x^k$ . Explain why  $S(x)/(1-x)$  is the generating function for the sums of squares. That is, the coefficient of  $x^n$  in the series for  $S(x)/(1-x)$  is  $\sum_{k=1}^n k^2$ .

(c) Use the fact that

$$S(x) = \frac{x(1+x)}{(1-x)^3},$$

and the previous part to prove that

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}.$$

(d) (Optional) How about a formula for the sum of cubes?

**Problem 2.** We are interested in generating functions for the number of different ways to compose a bag of  $n$  donuts subject to various restrictions. For each of the restrictions in (a)-(e) below, find a closed form for the corresponding generating function.

- (a) All the donuts are chocolate and there are at least 3.
- (b) All the donuts are glazed and there are at most 2.
- (c) All the donuts are coconut and there are exactly 2 or there are none.
- (d) All the donuts are plain and their number is a multiple of 4.
- (e) The donuts must be chocolate, glazed, coconut, or plain and:
- there must be at least 3 chocolate donuts, and
  - there must be at most 2 glazed, and
  - there must be exactly 0 or 2 coconut, and
  - there must be a multiple of 4 plain.
- (f) Find a closed form for the number of ways to select  $n$  donuts subject to the constraints of the previous part.

## Appendix

### Products of Series

Let

$$A(x) = \sum_{n=0}^{\infty} a_n x^n, \quad B(x) = \sum_{n=0}^{\infty} b_n x^n, \quad C(x) = A(x) \cdot B(x) = \sum_{n=0}^{\infty} c_n x^n.$$

Then

$$c_n = a_0 b_n + a_1 b_{n-1} + a_2 b_{n-2} + \cdots + a_n b_0.$$