

6	9	13	7
12	10	5	
3	1	4	14
15	8	11	2

Recursive Data



6	9	13	7
12	10	5	
3	1	4	14
15	8	11	2

Recursive Definitions

Define something in terms of a simpler version of the same thing:

Base case(s) that don't depend on anything else.

Constructor case(s) that depend on simpler cases.



6	9	13	7
12	10	5	
3	1	4	14
15	8	11	2

Example Definition: set E

Define set $E \subseteq \mathbb{Z}$,
recursively:

- **Base case:** $0 \in E$
- **Constructor cases:**

If $n \in E$, then

1. $n + 2 \in E$, if $n \geq 0$;
2. $-n \in E$, if $n > 0$.



6	9	13	7
12	10	5	
3	1	4	14
15	8	11	2

Example Definition: set E

1. $n \in E$ and $n \geq 0$, then $n + 2 \in E$:

$0, 0+2, (0+2)+2, ((0+2)+2)+2$

$0, 2, 4, 6, \dots$

2. $n \in E$ and $n > 0$, then $-n \in E$

$-2, -4, -6, \dots$

all even numbers



6	9	13	7
12	10	5	
3	1	4	14
15	8	11	2

Recursive Def: Extremal Clause

So, E contains the even integers
Anything Else? **No!**



Albert R Meyer, February 29, 2012

lec 5M.5

6	9	13	7
12	10	5	
3	1	4	14
15	8	11	2

Recursive Def: Extremal Clause

So, E contains the even integers

- $0 \in E$
- If $n \in E$ and $n \geq 0$, then $n+2 \in E$
- If $n \in E$ and $n > 0$, then $-n \in E$
- **That's All!**

Extremal Clause

(Implicit part of definition)



Albert R Meyer, February 29, 2012

lec 5M.6

6	9	13	7
12	10	5	
3	1	4	14
15	8	11	2

Example Definition: set E

So E is exactly
the Even Integers



Albert R Meyer, February 29, 2012

lec 5M.7

6	9	13	7
12	10	5	
3	1	4	14
15	8	11	2

Matched Paren Strings, M

set of strings, $M \subseteq \{], [\}^*$

- Base: $\lambda \in M$,
(the empty string)
- Constructor:

If $s, t \in M$, then

$$[s]t \in M$$



Albert R Meyer, February 29, 2012

lec 5M.9



Matched Paren Strings M

strings $[s]t \in M$

$[]$	$s = \lambda$	$t = \lambda$
$[][]$	$s = []$	$t = \lambda$
$[][][]$	$s = \lambda$	$t = []$
$[][]] []$	$s = []$	$t = []$
$[[[]]]$	$s = [[]]$	$t = \lambda$
\vdots	\vdots	\vdots



Albert R Meyer, February 29, 2012

7W.10



not in M

strings starting with $]$

are **not** in M because

- λ does not start with $]$
- $[s]t$ does not start with $]$

and everything in M arises in one of these two ways



Albert R Meyer, February 29, 2012

7W.11



The 18.01 Functions, $F18$

The set $F18$ of functions on \mathbb{R} :

$\text{Id}_{\mathbb{R}}$, constant functions, and $\sin x$ are in $F18$.

if $f, g \in F18$, then

- $f + g$, $f \cdot g$, 2^f ,
- the inverse, $f^{(-1)}$, of f , and
- $f \circ g$ (the composition of f and g) are in $F18$.



Albert R Meyer, February 29, 2012

lec 5M.13



The 18.01 Functions, $F18$

Some functions in $F18$:

$$-x = (-1) \cdot x$$

$$\sqrt{x} = (x^2)^{(-1)} \text{ ---inverse}$$

$$\cos x = (1 - (\sin x \cdot \sin x))^{1/2}$$

$$\ln x = (2^{\log x})^{(-1)}$$



Albert R Meyer, February 29, 2012

lec 5M.14

MIT OpenCourseWare
<http://ocw.mit.edu>

6.042J / 18.062J Mathematics for Computer Science
Spring 2015

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.