

MIT OpenCourseWare
<http://ocw.mit.edu>

6.641 Electromagnetic Fields, Forces, and Motion, Spring 2005

Please use the following citation format:

Markus Zahn, *6.641 Electromagnetic Fields, Forces, and Motion, Spring 2005*. (Massachusetts Institute of Technology: MIT OpenCourseWare).
<http://ocw.mit.edu> (accessed MM DD, YYYY). License: Creative Commons Attribution-Noncommercial-Share Alike.

Note: Please use the actual date you accessed this material in your citation.

For more information about citing these materials or our Terms of Use, visit:
<http://ocw.mit.edu/terms>

Massachusetts Institute of Technology
 Department of Electrical Engineering and Computer Science
 6.641 Electromagnetic Fields, Forces, and Motion

Problem Set #6
 Spring Term 2005

Issued: 3/15/05
 Due: 3/31/05

Suggested Reading Assignment: Zahn – Chapter 4.1-4.4.3; 5.7.2

Suggested Video Viewing: H/M Demos 10.7.1, 7.5.1, 7.5.2

[http://web.mit.edu/6.013_book/www/VideoDemo.html] and Demos: 10.0.1, 8.5.1, 10.2.1

Message: Right after the Spring Break we will be starting the text books *Electromechanical Dynamics, Part I: Discrete Systems and Part II: Fields, Forces, Motion* by H.H. Woodson and J.R. Melcher. Lecture notes will be handed out for all lectures, so it is not required for you to obtain these books.


Problem 6.1

The potential on an infinitely long cylinder of radius a is constrained to be

$$\Phi(r = a, \phi) = V_o \cos n\phi$$

The cylinder has permittivity ϵ and is surrounded by free space.

- (a) Find the potential and electric field everywhere inside and outside the cylinder.
- (b) What is the surface charge distribution on the cylindrical surface at $r=a$?
- (c)



Diagrams from *Electromagnetic Field Theory: A Problem Solving Approach*, by Markus Zahn, Robert E. Krieger Publishing Company, 1987. Used with permission.

The potential is now changed so that it is constant on each half of the cylinder:

$$\Phi(r = a, \phi) = \begin{cases} V_o / 2, & -\frac{\pi}{2} < \phi < \frac{\pi}{2} \\ -V_o / 2, & \frac{\pi}{2} < \phi < \frac{3\pi}{2} \end{cases}$$

Write this square wave of potential as a Fourier series.

- (d) Use the results of (a) and (c) to find the potential distribution due to this square wave of potential.

Problem 6.2


We wish to find solutions to Laplace's equation in cylindrical coordinates that have azimuthal symmetry so that the potential is independent of the angle ϕ , $\Phi(r, z)$.

- (a) Guess a product solution of the form

$$\Phi(r, z) = R(r)Z(z)$$

and find the zero separation constant solutions only. For these solutions of $R(r)$ and $Z(z)$ write down the general form for the potential.

- (b)



An Ohmic conducting cylinder with conductivity σ , permittivity ϵ , radius a , and height ℓ is surrounded by free space and a grounded perfectly conducting cylindrical can of radius b open at one end. A voltage V_0 is applied at the top of the resistive cylinder.

What are the boundary conditions on the potential at $z=0$, at $z=\ell$ for $r < a$, and at $r=b$? What does

$$[\Phi(r = a_+, z = \ell) - \Phi(r = a_-, z = \ell)]$$

equal?

From *Electromagnetic Field Theory: A Problem Solving Approach*, by Markus Zahn, Robert E. Krieger Publishing Company, 1987. Used with permission.

- (c) Use the general solutions from part (a) in each region for $0 \leq r \leq a$ and $a \leq r \leq b$ together with the boundary conditions in part (b) to find the potential distribution everywhere for $0 < r < b$ and $0 < z < \ell$.
- (d) What is the electric field distribution?
- (e) What is the free surface charge distribution and total charge on the interface at $r=a$?

- (f) What is the equation of the equipotential line and of the electric field line in the free space region that passes through the coordinate (r_0, z_0) where $a < r_0 < b$ and $0 < z < \ell$. Prove that the equipotential line and the electric field line cross perpendicularly.


Hint: $\frac{dr}{dz} = \frac{E_r}{E_z}$; $\int r \ln r dr = \frac{r^2}{2} \ln r - \frac{r^2}{4}$

Problem 6.3

An electric dipole having a z-directed moment $p(t)$ is situated at the origin and at the center of a spherical cavity of free space having a radius R in a material having uniform ϵ and σ . When $t < 0$, $p = 0$ and there is no free charge anywhere. The dipole is a step function of time, instantaneously assuming a moment p_0 when $t = 0$.

- An instant after the dipole is established, what is the distribution of Φ =inside and outside the cavity?
- Long after the electric dipole is turned on and the fields have reached a steady state, what is the distribution of Φ ?
- Determine $\Phi(r, \theta, t)$.
- What is the free surface charge density at $r = R$?


Problem 6.4



A sphere of radius R is permanently magnetized as $\bar{M} = M_0 \bar{i}_z$ and is surrounded by free space with magnetic permeability μ_0 .

Find the magnetic scalar potential χ and the magnetic field \bar{H} everywhere, where $\bar{H} = -\nabla \chi$.

Problem 6.5



A dielectric cylinder of infinite length and radius a is permanently polarized as $\bar{P} = P_0 \bar{i}_y$ and is surrounded by free space with dielectric permittivity ϵ_0 . There is no free surface charge at $r = a$.

- (a) What is the polarization surface charge distribution at $r = a$?

Hint: $\bar{i}_y = \sin \phi \bar{i}_r + \cos \phi \bar{i}_\phi$

- (b) What are the electric scalar potential $\Phi(r, \phi)$ and electric field $\bar{E} = -\nabla \Phi$.