

3.320: Lecture 15 (Mar 31 2005)

FIRST-PRINCIPLES MOLECULAR DYNAMICS

...and let us, as nature directs,
begin first with first principles.

Aristotle (Poetics, I)

Simulated Annealing




Figure by MIT OCW.

Micro- to macro- : diffusion coefficient

- From Fick to Einstein:

$$\boxed{\frac{\partial c(r,t)}{\partial t} = D \nabla^2 c(r,t)}$$

$$c(r,t)$$
$$\vec{j} = -D \nabla c$$
$$\frac{\partial c}{\partial t} + \vec{D} \cdot \vec{\nabla} c = 0$$

$$\frac{\partial}{\partial t} \int d\vec{r} r^2 c(r,t) = D \int d\vec{r} r^2 \nabla^2 c(r,t)$$



$$\frac{\partial}{\partial t} \underbrace{\langle r^2(t) \rangle}_{\uparrow} = 2dD$$

Mean Square Displacements

$$\langle \Delta r(t)^2 \rangle = \frac{1}{N} \sum_{i=1}^N \Delta r_i(t)^2$$

Mean Square Displacements

Liquid



Velocity Autocorrelation Function

$$\langle \Delta x(t)^2 \rangle = \frac{1}{N} \sum_{i=1}^N \Delta x_i(t)^2$$

$$\Delta x(t) = \int_0^t dt' v_x(t')$$


$$\langle \Delta x(t)^2 \rangle = \left\langle \left(\int_0^t dt' v_x(t') \right)^2 \right\rangle = \int_0^t dt' \int_0^t dt'' \langle v_x(t') v_x(t'') \rangle$$

$$= 2 \int_0^t dt' \int_0^{t'} dt'' \langle v_x(t') v_x(t'') \rangle$$



Green-Kubo relations

$$2D = \lim_{t \rightarrow \infty} \frac{\partial \langle x^2(t) \rangle}{\partial t} = \lim_{t' \rightarrow \infty} 2 \int_0^{t'} dt'' \langle v_x(t') v_x(t'') \rangle$$

$$\langle v_x(t') v_x(t'') \rangle = \langle v_x(t' - t'') v_x(0) \rangle$$




Velocity Autocorrelation Function



More Green-Kubo

- Other transport coefficients:
 - Shear viscosity, from the stress
 - Electrical conductivity, from the charge current
 - IR adsorption, from the polarization



Dynamics, Lagrangian style

- First construct $L = T - V$
 - Then, the equations of motion are given by
- $$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_j} \right) - \frac{\partial L}{\partial q_j} = 0$$
- (the dot is a time derivative)

- Why ? We can use generalized coordinates.
Also, we only need to think at the two scalar functions T and V

Newton's second law, too

- 1-d, 1 particle: $T=1/2 mv^2$, $V=V(x)$

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_j} \right) - \frac{\partial L}{\partial q_j} = 0$$
$$\frac{d}{dt} \left(\frac{\partial \left(\frac{1}{2} m \dot{x}^2 \right)}{\partial \dot{x}} \right) + \frac{\partial V}{\partial x} = 0 \quad \xrightarrow{\text{green arrow}} \quad \frac{d}{dt} (m \dot{x}) = - \frac{\partial V}{\partial x}$$

Hamiltonian

- We could use it to derive Hamiltonian dynamics (twice the number of differential equations, but all first order). We introduce a Legendre transformation

$$p_i = \frac{\partial L}{\partial \dot{q}_i} \quad H(q, p, t) = \sum_i \dot{q}_i p_i - L(q, \dot{q}, t)$$

$$\dot{q}_i = \frac{\partial H}{\partial p_i} \quad - \dot{p}_i = \frac{\partial H}{\partial q_i}$$


$$H = T + V$$

Thermostats, barostats...

- We might want to sample a constant-temperature ensemble, or constants pressure
 - Stochastic approach
 - Extended system
 - Constraint method


Nose' extended Lagrangian

$$L_{NOSE} = \sum_i \frac{1}{2} m_i s^2 \dot{r}_i^2 - V + \frac{1}{2} Q \dot{s}^2 - \frac{(3N+1)}{\beta} \ln s$$



Ergodicity issues

Very harmonic solids (e.g. 1 harmonic oscillator !)



Classical MD Bibliography



- Allen and Tildesley, *Computer Simulations of Liquids* (Oxford)
- Frenkel and Smit, *Understanding Molecular Simulations* (Academic)
- Ercolessi, *A Molecular Dynamics Primer* (<http://www.fisica.uniud.it/~ercolessi/md>)

First-principles molecular dynamics

Graph removed for copyright reasons.
Shows dramatic increase in number of citations per year of
“CP PRL 1985” and “AIMD” beginning around 1990.

Plane waves basis set

$$\vec{G}_i \cdot \vec{a}_j = 2\pi\delta_{ij}$$



It's really kinetic + potential

$$\hat{H} = -\frac{1}{2} \nabla^2 + V(\vec{r})$$

$$\psi_n(\vec{r}) = \sum_{\vec{G}} c_{\vec{G}}^n \exp(i \vec{G} \cdot \vec{r})$$


$$E = \sum_n \varepsilon_n = \sum_n \langle \psi_n | \hat{H} | \psi_n \rangle$$

Kinetic energy

$$E_{kin} = \sum_n \left\langle \psi_n \left| -\frac{1}{2} \nabla^2 \right| \psi_n \right\rangle \quad \psi_n(\vec{r}) = \sum_{\vec{G}} c_{\vec{G}}^n \exp(i \vec{G} \cdot \vec{r})$$

$$\left\langle G \left| -\frac{1}{2} \nabla^2 \right| G' \right\rangle = \int dr \exp(-iGr) \left[-\frac{1}{2} \nabla^2 \right] \exp(iG'r) = \frac{1}{2} G^2 \delta_{G,G'}$$

$$E_{kin} = \sum_n \frac{1}{2} \sum_{\vec{G}} \|c_{\vec{G}}^n\|^2 G^2$$



Total energy (non-SCF)

$$E_{pot} = \sum_n \langle \psi_n | V(\vec{r}) | \psi_n \rangle$$


$$\psi_n(\vec{r}) = \sum_{\vec{G}} c_{\vec{G}}^n \exp(i \vec{G} \cdot \vec{r})$$



$$\langle G | V(r) | G' \rangle = \int dr \exp(-iGr) V(r) \exp(iG'r) = V(G - G')$$

$$E_{tot} = \sum_n \left(\frac{1}{2} \sum_{\vec{G}} \|c_{\vec{G}}^n\|^2 G^2 + \sum_{\vec{G}, \vec{G}'} c_{\vec{G}}^{n*} c_{\vec{G}'}^n V(\vec{G} - \vec{G}') \right)$$

Dynamical evolution of c's




$$E_{tot} = \sum_n \left(\frac{1}{2} \sum_{\vec{G}} \|c_{\vec{G}}^n\|^2 G^2 + \sum_{\vec{G}, \vec{G}'} c_{\vec{G}}^{n*} c_{\vec{G}'}^n V(\vec{G} - \vec{G}') \right)$$

We need the force

$$E = E[\{\psi_i\}] \xrightarrow{\text{[green arrow]}} F_i = -\frac{\delta E[\{\psi_i\}]}{\delta \psi_i}$$
$$= -\hat{H}\psi_i$$

Skiing down a valley



Conjugate-gradients minimization


Hellmann-Feynman theorem

$$\vec{F}_i = -\frac{dE}{d\vec{R}_i} = -\frac{d\langle \Psi | \hat{H} | \Psi \rangle}{d\vec{R}_i} = \\ = \langle \Psi | -\frac{d\hat{H}}{d\vec{R}_i} | \Psi \rangle = \langle \Psi | -\frac{d\hat{V}}{d\vec{R}_i} | \Psi \rangle$$

Proof of Hellmann-Feynman

Born-Oppenheimer Molecular Dynamics

$$m_i \ddot{\vec{R}}_i = \vec{F}_i = \left\langle \Psi \left| -\frac{d\hat{V}}{d\vec{R}_i} \right| \Psi \right\rangle$$



The extended Car-Parrinello Lagrangian

$$\mathcal{L}_{\text{CP}} = \underbrace{\sum_I \frac{1}{2} M_I \dot{\mathbf{R}}_I^2 + \sum_i \frac{1}{2} \mu_i \langle \dot{\psi}_i | \dot{\psi}_i \rangle}_{\text{kinetic energy}} - \underbrace{\langle \Psi_0 | \mathcal{H}_e | \Psi_0 \rangle}_{\text{potential energy}} + \underbrace{\text{constraints}}_{\text{orthonormality}}$$

Equations of motion

$$M_I \ddot{\mathbf{R}}_I(t) = -\frac{\partial}{\partial \mathbf{R}_I} \langle \Psi_0 | \mathcal{H}_e | \Psi_0 \rangle$$



$$\mu_i \ddot{\psi}_i(t) = -\frac{\delta}{\delta \psi_i^\star} \langle \Psi_0 | \mathcal{H}_e | \Psi_0 \rangle + \frac{\delta}{\delta \psi_i^\star} \{ constraints \}$$

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{\mathbf{R}}_I} = \frac{\partial \mathcal{L}}{\partial \mathbf{R}_I}$$

$$\frac{d}{dt} \frac{\delta \mathcal{L}}{\delta \dot{\psi}_i^\star} = \frac{\delta \mathcal{L}}{\delta \psi_i^\star}$$

Constant of Motion

$$\underbrace{\sum_I \frac{1}{2} M_I \dot{\mathbf{R}}_I^2 + \sum_i \frac{1}{2} \mu_i \langle \dot{\psi}_i | \dot{\psi}_i \rangle}_{\text{kinetic energy}} + \underbrace{\langle \Psi_0 | \mathcal{H}_e | \Psi_0 \rangle}_{\text{potential energy}}$$



Born-Oppenheimer vs Car-Parrinello

BO vs CP forces

Kolmogorov-Arnold-Moser invariant tori

Quantum MD Bibliography

- Payne, Teter, Allan, Arias, Joannopoulos, *Rev Mod Physics* 64, 1045 (1992).
- Marx, Hutter, "Ab Initio Molecular Dynamics: Theory and Implementation", in "Modern Methods and Algorithms of Quantum Chemistry" (p. 301-449), Editor: J. Grotendorst, (NIC, FZ Jülich 2000)
- <http://www.theochem.ruhr-uni-bochum.de/research/marx/cprev.en.html>