

MIT OpenCourseWare
<http://ocw.mit.edu>

18.01 Single Variable Calculus
Fall 2006

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

Lecture 11: Max/Min Problems

Example 1. $y = \frac{\ln x}{x}$ (same function as in last lecture)




Figure 1: Graph of $y = \frac{\ln x}{x}$.

- What is the maximum value? Answer: $y = \frac{1}{e}$.
- Where (or at what point) is the maximum achieved? Answer: $x = e$. (See Fig. 1.)

Beware: Some people will ask “What is the maximum?”. The answer is *not* e . You will get so used to finding the critical point $x = e$, the main calculus step, that you will forget to find the maximum value $y = \frac{1}{e}$. Both the critical point $x = e$ and critical value $y = \frac{1}{e}$ are important. Together, they form the point of the graph $(e, \frac{1}{e})$ where it turns around.

Example 2. Find the max and the min of the function in Fig. 2

Answer: If you’ve already graphed the function, it’s obvious where the maximum and minimum values are. The point is to find the maximum and minimum without sketching the whole graph.

Idea: Look for the max and min among the critical points and endpoints. You can see from Fig. 2 that we only need to compare the heights or y -values corresponding to endpoints and critical points. (Watch out for discontinuities!)




Figure 2: Search for max and min among critical points and endpoints

Example 3. Find the open-topped can with the least surface area enclosing a fixed volume, V .




Figure 3: Open-topped can.

1. Draw the picture.
2. Figure out what variables to use. (In this case, r , h , V and surface area, S .)
3. Figure out what the constraints are in the problem, and express them using a formula. In this example, the constraint is

$$V = \pi r^2 h = \text{constant}$$

We're also looking for the surface area. So we need the formula for that, too:

$$S = \pi r^2 + (2\pi r)h$$

Now, in symbols, the problem is to minimize S with V constant.

4. Use the constraint equation to express everything in terms of r (and the constant V).

$$h = \frac{V}{2\pi r}; \quad S = \pi r^2 + (2\pi r) \left(\frac{V}{\pi r^2} \right)$$

5. Find the critical points (solve $dS/dr = 0$), as well as the endpoints. S will achieve its max and min at one of these places.

$$\frac{dS}{dr} = 2\pi r - \frac{2V}{r^2} = 0 \implies \pi r^3 - V = 0 \implies r^3 = \frac{V}{\pi} \implies r = \left(\frac{V}{\pi} \right)^{1/3}$$

We're not done yet. We've still got to evaluate S at the endpoints: $r = 0$ and " $r = \infty$ ".

$$S = \pi r^2 + \frac{2V}{r}, \quad 0 \leq r < \infty$$

As $r \rightarrow 0$, the second term, $\frac{2}{r}$, goes to infinity, so $S \rightarrow \infty$. As $r \rightarrow \infty$, the first term πr^2 goes to infinity, so $S \rightarrow \infty$. Since $S = +\infty$ at each end, the minimum is achieved at the critical point $r = (V/\pi)^{1/3}$, not at either endpoint.




Figure 4: Graph of S

We're still not done. We want to find the minimum value of the surface area, S , and the values of h .

$$r = \left(\frac{V}{\pi} \right)^{1/3}; \quad h = \frac{V}{\pi r^2} = \frac{V}{\pi \left(\frac{V}{\pi} \right)^{2/3}} = \frac{V}{\pi} \left(\frac{V}{\pi} \right)^{-2/3} = \left(\frac{V}{\pi} \right)^{1/3}$$

$$S = \pi r^2 + 2\frac{V}{r} = \pi \left(\frac{V}{\pi} \right)^{2/3} + 2V \left(\frac{V}{\pi} \right)^{1/3} = 3\pi^{-1/3} V^{2/3}$$

Finally, another, often better, way of answering that question is to find the proportions of the can. In other words, what is $\frac{h}{r}$? Answer: $\frac{h}{r} = \frac{(V/\pi)^{1/3}}{(V/\pi)^{1/3}} = 1$.

Example 4. Consider a wire of length 1, cut into two pieces. Bend each piece into a square. We want to figure out where to cut the wire in order to enclose as much area in the two squares as possible.




Figure 5: Illustration for Example 5.

The first square will have sides of length $\frac{x}{4}$. Its area will be $\frac{x^2}{16}$. The second square will have sides of length $\frac{1-x}{4}$. Its area will be $(\frac{1-x}{4})^2$. The total area is then

$$A = \left(\frac{x}{4}\right)^2 + \left(\frac{1-x}{4}\right)^2$$

$$A' = \frac{2x}{16} + \frac{2(1-x)}{16}(-1) = \frac{x}{8} - \frac{1}{8} + \frac{x}{8} = 0 \implies 2x - 1 = 0 \implies x = \frac{1}{2}$$

So, one extreme value of the area is

$$A = \left(\frac{1/2}{4}\right)^2 + \left(\frac{1/2}{4}\right)^2 = \frac{1}{32}$$

We're not done yet, though. We still need to check the endpoints! At $x = 0$,

$$A = 0^2 + \left(\frac{1-0}{4}\right)^2 = \frac{1}{16}$$

At $x = 1$,

$$A = \left(\frac{1}{4}\right)^2 + 0^2 = \frac{1}{16}$$

By checking the endpoints in Fig. 6, we see that the *minimum* area was achieved at $x = \frac{1}{2}$. The maximum area is not achieved in $0 < x < 1$, but it is achieved at $x = 0$ or 1 . The maximum corresponds to using the whole length of wire for one square.




Figure 6: Graph of the area function.

Moral: Don't forget endpoints. If you only look at critical points you may find the worst answer, rather than the best one.