

MIT OpenCourseWare  
<http://ocw.mit.edu>

18.02 Multivariable Calculus  
Fall 2007

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

## 18.02 Practice Exam 2 B – Solutions


**Problem 1.** a)  $\nabla f = \langle 2xy^2 - 1, 2x^2y \rangle = \langle 3, 8 \rangle = 3\hat{i} - 8\hat{j}$ .

b)  $z - 2 = 3(x - 2) + 8(y - 1)$  or  $z = 3x + 8y - 12$ .

c)  $\Delta x = 1.9 - 2 = -1/10$  and  $\Delta y = 1.1 - 1 = 1/10$ . So  $z \simeq 2 + 3\Delta x + 8\Delta y = 2 - 3/10 + 8/10 = 2.5$

d)  $\left. \frac{df}{ds} \right|_{\hat{u}} = \nabla f \cdot \hat{u} = \langle 3, 8 \rangle \cdot \frac{\langle -1, 1 \rangle}{\sqrt{2}} = \frac{-3 + 8}{\sqrt{2}} = \frac{5}{\sqrt{2}}$

**Problem 2.**



**Problem 3.** a) 
$$\left. \begin{aligned} w_x = -6x - 4y + 16 = 0 &\Rightarrow -3x - 2y + 8 = 0 \\ w_y = -4x - 2y - 12 = 0 &\Rightarrow 4x + 2y + 12 = 0 \end{aligned} \right\} \Rightarrow \begin{cases} x = -20 \\ y = 34 \end{cases}$$

Therefore there is just one critical point at  $(-20, 34)$ . Since

$$w_{xx}w_{yy} - w_{xy}^2 = (-6)(-2) - (-4)^2 = 12 - 16 = -4 < 0,$$

the critical point is a saddle point.

b) There is no critical point in the first quadrant, hence the maximum must be at infinity or on the boundary of the first quadrant.

The boundary is composed of two half-lines:

- $x = 0$  and  $y \geq 0$  on which  $w = -y^2 - 12y$ . It has a maximum ( $w = 0$ ) at  $y = 0$ .
- $y = 0$  and  $x \geq 0$ , where  $w = -3x^2 + 16x$ . (The graph is a parabola pointing downwards).  
Maximum:  $w_x = -6x + 16 = 0 \Rightarrow x = 8/3$ . Hence  $w$  has a maximum at  $(8/3, 0)$  and  $w = -3(8/3)^2 + 16 \cdot 8/3 = 64/3 > 0$ .

We now check that the maximum of  $w$  is not at infinity:

- If  $y \geq 0$  and  $x \rightarrow +\infty$  then  $w \leq -3x^2 + 16x$ , which tends to  $-\infty$  as  $x \rightarrow +\infty$ .
- If  $0 \leq x \leq C$  and  $y \rightarrow +\infty$ , then  $w \leq -y^2 + 16C$ , which tends to  $-\infty$  as  $y \rightarrow +\infty$ .

We conclude that the maximum of  $w$  in the first quadrant is at  $(8/3, 0)$ .

**Problem 4.** a) 
$$\begin{cases} w_x = u_x w_u + v_x w_v = -\frac{y}{x^2} w_u + 2x w_v \\ w_y = u_y w_u + v_y w_v = \frac{1}{x} w_u + 2y w_v \end{cases}$$

b)  $xw_x + yw_y = x\left(-\frac{y}{x^2} w_u + 2x w_v\right) + y\left(\frac{1}{x} w_u + 2y w_v\right) = \left(-\frac{y}{x} + \frac{y}{x}\right) w_u + (2x^2 + 2y^2) w_v = 2vw_v$ .

c)  $xw_x + yw_y = 2vw_v = 2v \cdot 5v^4 = 10v^5$ .

**Problem 5.** a)  $f(x, y, z) = x$ ; the constraint is  $g(x, y, z) = x^4 + y^4 + z^4 + xy + yz + zx = 6$ . The Lagrange multiplier equation is:

$$\nabla f = \lambda \nabla g \quad \Leftrightarrow \quad \begin{cases} 1 &= \lambda(4x^3 + y + z) \\ 0 &= \lambda(4y^3 + x + z) \\ 0 &= \lambda(4z^3 + x + y) \end{cases}$$

b) The level surfaces of  $f$  and  $g$  are tangent at  $(x_0, y_0, z_0)$ , so they have the same tangent plane. The level surface of  $f$  is the plane  $x = x_0$ ; hence this is also the tangent plane to the surface  $g = 6$  at  $(x_0, y_0, z_0)$ .

Second method: at  $(x_0, y_0, z_0)$ , we have

$$\left. \begin{array}{l} 1 = \lambda g_x \\ 0 = \lambda g_y \\ 0 = \lambda g_z \end{array} \right\} \Rightarrow \lambda \neq 0 \text{ and } \langle g_x, g_y, g_z \rangle = \langle \frac{1}{\lambda}, 0, 0 \rangle.$$

So  $\langle \frac{1}{\lambda}, 0, 0 \rangle$  is perpendicular to the tangent plane at  $(x_0, y_0, z_0)$ ; the equation of the tangent plane is then  $\frac{1}{\lambda}(x - x_0) = 0$ , or equivalently  $x = x_0$ .

**Problem 6.**

a) Taking the total differential of  $x^2 + y^3 - z^4 = 1$ , we get:  $2x dx + 3y^2 dy - 4z^3 dz = 0$ . Similarly, from  $z^3 + zx + xy = 3$ , we get:  $(y + z) dx + x dy + (3z^2 + x) dz = 0$ .

b) At  $(1, 1, 1)$  we have:  $2 dx + 3 dy - 4 dz = 0$  and  $2 dx + dy + 4 dz = 0$ . We eliminate  $dz$  (by adding these two equations):  $4 dx + 4 dy = 0$ , so  $dy = -dx$ , and hence  $dy/dx = -1$ .