

18.02 Practice Exam 1A - Solutions

Problem 1.

a) $\overrightarrow{OQ} = \hat{i} + \hat{j} + \hat{k}$; $\overrightarrow{OR} = \frac{1}{2}\hat{i} + \hat{j} + \frac{1}{2}\hat{k}$.

b) $\cos \theta = \frac{\overrightarrow{OQ} \cdot \overrightarrow{OR}}{|\overrightarrow{OQ}| |\overrightarrow{OR}|} = \frac{\langle 1, 1, 1 \rangle \cdot \langle \frac{1}{2}, 1, \frac{1}{2} \rangle}{\sqrt{3} \sqrt{\frac{3}{2}}} = \frac{2\sqrt{2}}{3}$.

Problem 2.

Velocity: $\vec{V} = \frac{d\vec{R}}{dt} = \langle -3 \sin t, 3 \cos t, 1 \rangle$. Speed: $|\vec{V}| = \sqrt{9 \sin^2 t + 9 \cos^2 t + 1} = \sqrt{10}$.

Problem 3.

a) Minors: $\begin{bmatrix} 1 & 1 & 2 \\ -2 & -2 & -2 \\ -3 & -5 & -6 \end{bmatrix}$. Cofactors: $\begin{bmatrix} 1 & -1 & 2 \\ 2 & -2 & 2 \\ -3 & 5 & -6 \end{bmatrix}$. Inverse: $\frac{1}{2} \begin{bmatrix} 1 & \boxed{2} & \boxed{-3} \\ -1 & -2 & 5 \\ 2 & 2 & -6 \end{bmatrix}$.

b) $X = A^{-1}B = \begin{bmatrix} -3 \\ 4 \\ -4 \end{bmatrix}$.

Problem 4.

Q = top of the ladder: $\overrightarrow{OQ} = \langle 0, L \sin \theta \rangle$; R = bottom of the ladder: $\overrightarrow{OR} = \langle -L \cos \theta, 0 \rangle$.

Midpoint: $\overrightarrow{OP} = \frac{1}{2}(\overrightarrow{OQ} + \overrightarrow{OR}) = \langle -\frac{L}{2} \cos \theta, \frac{L}{2} \sin \theta \rangle$.

Parametric equations: $x = -\frac{L}{2} \cos \theta$, $y = \frac{L}{2} \sin \theta$.

Problem 5.

a) $\overrightarrow{P_0P_1} \times \overrightarrow{P_0P_2} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & -1 & 1 \\ 0 & -2 & 1 \end{vmatrix} = \hat{i} + \hat{j} + 2\hat{k}$. Area = $\frac{1}{2} |\overrightarrow{P_0P_1} \times \overrightarrow{P_0P_2}| = \frac{1}{2}\sqrt{6}$.

b) Normal vector: $\overrightarrow{P_0P_1} \times \overrightarrow{P_0P_2} = \hat{i} + \hat{j} + 2\hat{k}$. Equation: $x + y + 2z = 3$.

c) Parametric equations for the line: $x = -1 + t$, $y = t$, $z = t$.

Substituting: $-1 + 4t = 3$, $t = 1$, intersection point $(0, 1, 1)$.

Problem 6.

a) $\frac{d}{dt}(\vec{R} \cdot \vec{R}) = \vec{V} \cdot \vec{R} + \vec{R} \cdot \vec{V} = 2\vec{R} \cdot \vec{V}$.

b) Assume $|\vec{R}|$ is constant: then $\frac{d}{dt}(\vec{R} \cdot \vec{R}) = 2\vec{R} \cdot \vec{V} = 0$, i.e. $\vec{R} \perp \vec{V}$.

c) $\vec{R} \cdot \vec{V} = 0$, so $\frac{d}{dt}(\vec{R} \cdot \vec{V}) = \vec{V} \cdot \vec{V} + \vec{R} \cdot \vec{A} = 0$. Therefore $\vec{R} \cdot \vec{A} = -|\vec{V}|^2$.