

## 18.02 Practice Exam 3 A – Solutions

**1.** a) The area of the triangle is 2, so  $\bar{y} = \frac{1}{2} \int_0^1 \int_{2y-2}^{2-2y} y \, dx \, dy$ .

b) By symmetry  $\bar{x} = 0$ .

**2.**  $\delta = |x| = |r \cos \theta|$ .  $I_0 = \iint_D r^2 \delta \, r \, dr \, d\theta = \int_0^{2\pi} \int_0^1 r^2 |r \cos \theta| \, r \, dr \, d\theta = 4 \int_0^{\pi/2} \int_0^1 r^4 \cos \theta \, r \, dr \, d\theta = 4 \int_0^{\pi/2} \frac{1}{5} \cos \theta \, d\theta = \frac{4}{5}$

**3.** a)  $N_x = 6x^2 + by^2$ ,  $M_y = ax^2 + 3y^2$ .  $N_x = M_y$  provided  $a = 6$  and  $b = 3$ .

b)  $f_x = 6x^2y + y^3 + 1 \implies f = 2x^3y + xy^3 + x + c(y)$ . Therefore,  $f_y = 2x^3 + 3xy^2 + c'(y)$ . Setting this equal to  $N$ , we have  $2x^3 + 3xy^2 + c'(y) = 2x^3 + 3xy^2 + 2$  so  $c'(y) = 2$  and  $c = 2y$ . So

$$f = 2x^3y + xy^3 + x + 2y \quad (+\text{constant}).$$

c)  $C$  starts at  $(1, 0)$  and ends at  $(-e^\pi, 0)$ , so  $\int_C \vec{F} \cdot d\vec{r} = f(-e^\pi, 0) - f(1, 0) = -e^{-\pi} - 1$ .

**4.**  $\int_C yx^3 \, dx + y^2 \, dy = \int_0^1 x^2 x^3 \, dx + (x^2)^2 (2x \, dx) = \int_0^1 3x^5 \, dx = 1/2$ .

**5.** a)  $\begin{vmatrix} u_x & u_y \\ v_x & v_y \end{vmatrix} = \begin{vmatrix} 2x/y & -x^2/y^2 \\ y & x \end{vmatrix} = 3x^2/y$ . Therefore,

$$dudv = (3x^2/y) \, dx \, dy = 3u \, dx \, dy \implies dx \, dy = \frac{1}{3u} \, dudv.$$

b)  $\int_2^4 \int_1^5 \frac{1}{3u} \, dudv = \int_2^4 \frac{1}{3} \ln 5 \, dv = \frac{2}{3} \ln 5$ .

**6.** a)  $\oint_C M \, dx = \iint_R -M_y \, dA$ .

b) We want  $M$  such that  $-M_y = (x+y)^2$ . Use  $M = -\frac{1}{3}(x+y)^3$ .

**7.** a)  $\operatorname{div} \vec{F} = 2y$ , so  $\iint_R 2y \, dA = \int_0^1 \int_0^{x^3} 2y \, dy \, dx = \int_0^1 x^6 \, dx = \frac{1}{7}$ .

b) For the flux through  $C_1$ ,  $\hat{\mathbf{n}} = -\hat{\mathbf{j}}$  implies  $\vec{F} \cdot \hat{\mathbf{n}} = -(1+y^2) = -1$  where  $y = 0$ . The length of  $C_1$  is 1, so the total flux through  $C_1$  is  $-1$ .

The flux through  $C_2$  is zero because  $\hat{\mathbf{n}} = \hat{\mathbf{i}}$  and  $\vec{F} \perp \hat{\mathbf{i}}$ .

c)  $\int_{C_3} \vec{F} \cdot \hat{\mathbf{n}} \, ds = \iint_R \operatorname{div} \vec{F} \, dA - \int_{C_1} \vec{F} \cdot \hat{\mathbf{n}} \, ds - \int_{C_2} \vec{F} \cdot \hat{\mathbf{n}} \, ds = \frac{1}{7} - (-1) - 0 = \frac{8}{7}$ .

MIT OpenCourseWare  
<http://ocw.mit.edu>

18.02SC Multivariable Calculus

Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.