

18.06 Linear Algebra, Fall 2011
Recitation Transcript – Solving $Ax=b$

MARTINA BALAGOVIC: Hi. Welcome.

Today's problem is about finding solutions of this non-homogeneous linear system, x minus $2y$ minus $2z$ equals b_1 , $2x$ minus $5y$ minus $4z$ equals b_2 , and $4x$ minus $9y$ minus $8z$ equals b_3 . And as you can see, the system doesn't only have numbers and unknowns, it also has parameters, b_1 , b_2 , and b_3 , and the solution will depend on these parameters, but also the existence of the solution will depend on these parameters. And we're asked to find a solution and find when it exists, depending on the values of b_1 , b_2 , and b_3 .

So now you should pause the video, solve the problem, and come back and compare your solution with mine.

And we're back. Let's try it. Let's start by solving this system as though b_1 , b_2 , and b_3 were numbers.

So we write the matrix of the system, which is 1 minus 2 minus 2 , b_1 , and then 2 minus 5 minus 4 , b_2 , and 4 minus 9 minus 8 , b_3 . And we do elimination. So we multiply the first row by minus 2 and add it to the second row. And we multiply it by minus 4 and add it to the third row. And we get 1 minus 2 minus 2 b_1 , 0 , 4 minus 5 is minus 1 , 4 minus 4 is 0 , and minus 2 times b_1 plus b_2 . And here we get 0 , 8 minus 9 is minus 1 , and 8 minus 8 is 0 . Finally on the right-hand side, minus $4b_1$ plus b_3 . And you can already see that something's going to happen here. But let's do one more step.

So eliminating further we get 1 minus 2 minus 2 b_1 , 0 minus 1 0 minus $2b_1$ plus b_2 . And in the last row we replace it with the last row minus the second row, and we get 0 0 0 minus $4b_1$ plus 2 -- so minus minus $2b_2$ is minus $2b_1$ minus b_2 and plus b_2 . I hope I did this right.

So now let's think of it as a system again. The last equation says 0 equals this expression in b_1 , b_2 , and b_3 . So this is something to note down. If minus $2b_1$ minus b_2 plus b_3 is some number that's not 0 , then the last equation is going to say 0 equals non- 0 . It's never going to be satisfied, and the entire system is never going to have a solution. So in this case, we have no solutions.

If this is equal to 0, so $-2b_1 - b_2 + b_3$ is equal to 0, then let's do one more step on this matrix here. Let's turn this number into 1 by multiplying this row by negative 1. And let's use it to eliminate this number here as well.

So in this case, we get-- let me write it from the last row, which now becomes $0\ 0\ 0$ equals 0, which is fine. The second row becomes $0\ 1\ 0\ 2b_1 - b_2$. And the first one to get rid of this minus 2, we multiply this row by negative 2 and add it to the first one. We get $1\ 0\ -2$, and here we get $b_1 + 4b_1$, which is $5b_1 - 2b_2$.

The reason why we did it was to get the identity matrix here. And now let's solve this.

These two columns corresponding to variables x and y have pivots in them. So these are the pivot variables. This column here has no pivot in it, so it's a free variable. And now we're going to calculate the solutions, but by picking particular values for z , and then calculating the values for x and y , which we have two kinds of solution. One kind is the particular solution. So this one solves $Ax = b$. There's only one of them. And we get it by setting the free variable equal to 0.

Setting the free variable equal to 0, we get, well this is equal to 0. The second equation says y equals this thing here, so $2b_1 - b_2$. And the first equation says $x - 2 \times 0$ equals this expression here. So $5b_1 - 2b_2$. That's our particular solution.

The next kind is the special solution. So remember, those solve $Ax = 0$. There's as many of them as there are free variables. In our case, there's only one. And we get it by setting all free variables equal to 0, except one equal to 1. And do it for every free variable. So in our case there's only one free variable and we set z equal to 1.

The solution that we get in this case, and remember we're solving $Ax = 0$ -- we don't care about the right-hand side anymore-- so z is 1. This second equation says $y = 0$, and the first equation says $x - 2 \times 1 = 0$. In other words, $x = 2$. So the special solution is $2\ 0\ 1$. And now all solutions are of the form x equals a particular solution plus any multiple of the special solution.

Let me recap. In case this particular combination of parameters is not 0, there's no solutions. In case this particular combination of parameters is equal to 0, there are solutions, there are as many of them as there are real numbers c , and they're all of this form for these two vectors. And that's all I wanted to say today.

MIT OpenCourseWare
<http://ocw.mit.edu>

18.06SC Linear Algebra
Fall 2011

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.