

Exercises on eigenvalues and eigenvectors

Problem 21.1: (6.1 #19. *Introduction to Linear Algebra: Strang*) A three by three matrix B is known to have eigenvalues 0, 1 and 2. This information is enough to find three of these (give the answers where possible):

- a) The rank of B
- b) The determinant of $B^T B$
- c) The eigenvalues of $B^T B$
- d) The eigenvalues of $(B^2 + I)^{-1}$

Solution:

- a) B has 0 as an eigenvalue and is therefore singular (not invertible). Since B is a three by three matrix, this means that its rank can be at most 2. Since B has two distinct nonzero eigenvalues, its rank is exactly 2.
- b) Since B is singular, $\det(B) = 0$. Thus $\det(B^T B) = \det(B^T) \det(B) = 0$.
- c) There is not enough information to find the eigenvalues of $B^T B$. For example:

$$\begin{aligned} \text{If } B = \begin{bmatrix} 0 & & \\ & 1 & \\ & & 2 \end{bmatrix} & \text{ then } B^T B = \begin{bmatrix} 0 & & \\ & 1 & \\ & & 4 \end{bmatrix}. \\ \text{If } B = \begin{bmatrix} 0 & 1 & \\ & 1 & \\ & & 2 \end{bmatrix} & \text{ then } B^T B = \begin{bmatrix} 0 & & \\ & 2 & \\ & & 4 \end{bmatrix}. \end{aligned}$$

- d) If $p(t)$ is a polynomial and if \mathbf{x} is an eigenvector of A with eigenvalue λ , then

$$p(A)\mathbf{x} = p(\lambda)\mathbf{x}.$$

We also know that if λ is an eigenvalue of A then $1/\lambda$ is an eigenvalue of A^{-1} . Hence the eigenvalues of $(B^2 + I)^{-1}$ are $\frac{1}{0^2+1}$, $\frac{1}{1^2+1}$ and $\frac{1}{2^2+1}$, or **1, 1/2 and 1/5.**

Problem 21.2: (6.1 #29.) Find the eigenvalues of $A, B,$ and C when

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}, B = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 2 & 0 \\ 3 & 0 & 0 \end{bmatrix} \text{ and } C = \begin{bmatrix} 2 & 2 & 2 \\ 2 & 2 & 2 \\ 2 & 2 & 2 \end{bmatrix}.$$

Solution: Since the eigenvalues of a triangular matrix are its diagonal entries, the eigenvalues of A are 1, 4, and 6. For B we have:

$$\begin{aligned} \det(B - \lambda I) &= (-\lambda)(2 - \lambda)(-\lambda) - 3(2 - \lambda) \\ &= (\lambda^2 - 3)(2 - \lambda). \end{aligned}$$

Hence the eigenvalues of B are $\pm\sqrt{3}$ and 2. Finally, for C we have:

$$\begin{aligned} \det(C - \lambda I) &= (2 - \lambda)[(2 - \lambda)^2 - 4] - 2[2(2 - \lambda) - 4] + 2[4 - 2(2 - \lambda)] \\ &= \lambda^3 - 6\lambda^2 = \lambda^2(\lambda - 6). \end{aligned}$$

The eigenvalues of C are 6, 0, and 0.

We can quickly check our answers by computing the determinants of A and B and by noting that C is singular.

MIT OpenCourseWare
<http://ocw.mit.edu>

18.06SC Linear Algebra
Fall 2011

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.