

## Exercises on orthogonal matrices and Gram-Schmidt

**Problem 17.1:** (4.4 #10.b *Introduction to Linear Algebra: Strang*)

Orthonormal vectors are automatically linearly independent.

Matrix Proof: Show that  $Q\mathbf{x} = \mathbf{0}$  implies  $\mathbf{x} = \mathbf{0}$ . Since  $Q$  may be rectangular, you can use  $Q^T$  but not  $Q^{-1}$ .

**Solution:** By definition,  $Q$  is a matrix whose columns are orthonormal, and so we know that  $Q^T Q = I$  (where  $Q$  may be rectangular). Then:

$$Q\mathbf{x} = \mathbf{0} \implies Q^T Q\mathbf{x} = Q^T \mathbf{0} \implies I\mathbf{x} = \mathbf{0} \implies \mathbf{x} = \mathbf{0}.$$

Thus the nullspace of  $Q$  is the zero vector, and so the columns of  $Q$  are linearly independent. There are no non-zero linear combinations of the columns that equal the zero vector. Thus, orthonormal vectors are automatically linearly independent.

**Problem 17.2:** (4.4 #18) Given the vectors  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  listed below, use the Gram-Schmidt process to find orthogonal vectors  $\mathbf{A}$ ,  $\mathbf{B}$ , and  $\mathbf{C}$  that span the same space.

$$\mathbf{a} = (1, -1, 0, 0), \mathbf{b} = (0, 1, -1, 0), \mathbf{c} = (0, 0, 1, -1).$$

Show that  $\{\mathbf{A}, \mathbf{B}, \mathbf{C}\}$  and  $\{\mathbf{a}, \mathbf{b}, \mathbf{c}\}$  are bases for the space of vectors perpendicular to  $\mathbf{d} = (1, 1, 1, 1)$ .

**Solution:** We apply Gram-Schmidt to  $\mathbf{a}, \mathbf{b}, \mathbf{c}$ . First, we set

$$\mathbf{A} = \mathbf{a} = (1, -1, 0, 0).$$

Next we find  $\mathbf{B}$ :

$$\mathbf{B} = \mathbf{b} - \frac{\mathbf{A}^T \mathbf{b}}{\mathbf{A}^T \mathbf{A}} \mathbf{A} = (0, 1, -1, 0) + \frac{1}{2}(1, -1, 0, 0) = \left(\frac{1}{2}, \frac{1}{2}, -1, 0\right).$$

And then we find  $\mathbf{C}$ :

$$\mathbf{C} = \mathbf{c} - \frac{\mathbf{A}^T \mathbf{c}}{\mathbf{A}^T \mathbf{A}} \mathbf{A} - \frac{\mathbf{B}^T \mathbf{c}}{\mathbf{B}^T \mathbf{B}} \mathbf{B} = (0, 0, 1, -1) + \frac{2}{3} \left(\frac{1}{2}, \frac{1}{2}, -1, 0\right) = \left(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, -1\right).$$

We know from the first problem that the elements of the set  $\{\mathbf{A}, \mathbf{B}, \mathbf{C}\}$  are linearly independent, and each vector is orthogonal to  $(1,1,1,1)$ . The space of vectors perpendicular to  $\mathbf{d}$  is three dimensional (since the row space of  $(1,1,1,1)$  is one-dimensional, and the number of dimensions of the row space added to the number of dimensions of the nullspace add to 4). Therefore  $\{\mathbf{A}, \mathbf{B}, \mathbf{C}\}$  forms a basis for the space of vectors perpendicular to  $\mathbf{d}$ .

Similarly,  $\{\mathbf{a}, \mathbf{b}, \mathbf{c}\}$  is a basis for the space of vectors perpendicular to  $\mathbf{d}$  because the vectors are linearly independent, orthogonal to  $(1,1,1,1)$ , and because there are three of them.

MIT OpenCourseWare  
<http://ocw.mit.edu>

18.06SC Linear Algebra  
Fall 2011

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.