

## Exercises on orthogonal vectors and subspaces

**Problem 16.1:** (4.1 #7. *Introduction to Linear Algebra: Strang*) For every system of  $m$  equations with no solution, there are numbers  $y_1, \dots, y_m$  that multiply the equations so they add up to  $0 = 1$ . This is called *Fredholm's Alternative*:

Exactly one of these problems has a solution:  
 $A\mathbf{x} = \mathbf{b}$  OR  $A^T\mathbf{y} = \mathbf{0}$  with  $\mathbf{y}^T\mathbf{b} = 1$ .

If  $\mathbf{b}$  is not in the column space of  $A$  it is not orthogonal to the nullspace of  $A^T$ . Multiply the equations  $x_1 - x_2 = 1$ ,  $x_2 - x_3 = 1$  and  $x_1 - x_3 = 1$  by numbers  $y_1, y_2$  and  $y_3$  chosen so that the equations add up to  $0 = 1$ .

**Problem 16.2:** (4.1#32.) Suppose I give you four nonzero vectors  $\mathbf{r}$ ,  $\mathbf{n}$ ,  $\mathbf{c}$  and  $\mathbf{l}$  in  $\mathbb{R}^2$ .

- a) What are the conditions for those to be bases for the four fundamental subspaces  $C(A^T)$ ,  $N(A)$ ,  $C(A)$ , and  $N(A^T)$  of a 2 by 2 matrix?
- b) What is one possible matrix  $A$ ?

MIT OpenCourseWare  
<http://ocw.mit.edu>

18.06SC Linear Algebra  
Fall 2011

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.