

MIT OpenCourseWare  
<http://ocw.mit.edu>

18.085 Computational Science and Engineering I  
Fall 2008

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

Your PRINTED name is: \_\_\_\_\_

Grading 1

2

3

\_\_\_\_\_

- 1) (30 pts.) (a) Suppose  $f(x)$  is a *periodic* function:

$$f(x) = \begin{cases} 0 & \text{for } -\pi < x < 0 \\ e^{-x} & \text{for } 0 \leq x \leq \pi \\ f(x + 2\pi n) & \text{for every integer } n \end{cases}$$

Find the coefficients  $c_k$  in the complex Fourier series  $f(x) = \sum c_k e^{ikx}$ .

What is  $c_0$ ? What is  $\sum_{-\infty}^{\infty} |c_k|^2$ ?

- (b) Draw a graph of  $f(x)$  from  $-2\pi$  to  $2\pi$ . Also draw a careful graph of  $df/dx$ . How quickly do the coefficients of  $f(x)$  decay as  $k \rightarrow \infty$  and why?
- (c) Find the Fourier coefficients  $d_k$  of  $df/dx$ . Do they approach a constant (or what pattern do they approach) as  $k \rightarrow \infty$ ? Explain the pattern from your graphs.



2) (33 pts.) (a) Can you complete this 4-step MATLAB code to compute the cyclic convolution  $f \circledast g = h$ ? I suggest `fhat`, `ghat`, `hhat` for their transforms.

1. `fhat = fft(f)`

2.

3. `hhat =`

4. `h =`

(It is equally possible to start with the inverse discrete transform `ifft`. The only difference will be a factor of  $N$  somewhere, which I forgive! If you don't know MATLAB notation for commands 2, 3, 4 you can use words. MATLAB's `fft(f)` and `ifft(f)` automatically determine the length of  $f$ .)

(b) Suppose each of your quiz grades is a random variable (don't know how I thought of this). The probability of grade  $j$  on each quiz ( $j = 0, \dots, 100$ ) is  $p_j$ . The "generating function" for that quiz is  $P(z) = \sum p_j z^j$ . What is the probability  $s_k$  that the sum of your grades on 2 quizzes is  $k$ ? Give a nice formula for  $S(z) = \sum s_k z^k$ .

(c) The chance of grade  $j = (70, 80, 90, 100)$  on one quiz is  $p = (.3, .4, .2, .1)$ . What is the expected value (mean  $m$ ) for the grade on that quiz? Show that this quiz average  $m$  agrees with  $dP/dz$  at  $z = 1$ . What are the probabilities  $s_k$  for the sum of two grades? Give numbers or a MATLAB code for the  $s_k$ .



- 3) (37 pts.) (a) The hat function  $H(x) = 1 - |x|$  for  $-1 \leq x \leq 1$  has height 1 and area 1 and integral transform  $\widehat{H}(k) = (2 - 2 \cos k)/k^2$ . Find the transform  $\widehat{R}(k)$  of the roof function  $R(x)$ :

$$R(x) = \mathbf{box} + \mathbf{hat} = 2 - |x| \quad \text{for } -1 \leq x \leq 1, \quad 0 \text{ else.}$$

- (b) What is the value of  $\widehat{R}(k)$  at  $k = 0$  and how does this connect to the graph of the roof?
- (c) Suppose  $R(x)$  is the response of a sensor to a point source  $\delta(x)$  at  $x = 0$ . The sensor is shift-invariant (shifted response when source is shifted). The output  $F$  from a distributed source  $U(x)$  is the convolution  $F = R * U$ . Describe how to find  $U(x)$  if you know  $F(x)$ .
- (d) There could be a difficulty with your solution method in part (c). That would arise if \_\_\_\_\_ = 0. For 1 point, does this difficulty appear in this example?

XXX