

# Classical Mechanics III (8.09) Fall 2014

## Assignment 5

Massachusetts Institute of Technology  
Physics Department  
Mon. October 6, 2014

*Due Tues. October 14, 2014  
6:00pm*

### Announcements

This week we continue to study Canonical Transformations.

- Due to the Columbus day holiday I have made this assignment due on Tuesday Oct. 14 rather than on the Monday. Note however that your next assignment (#6) will be posted on Monday Oct.13 and due on Monday Oct. 20 as usual.

### Reading Assignment

- The reading for Canonical Transformations is **Goldstein** Ch.9 sections 9.1-9.7. (We will not discuss active infinitesimal canonical transformations with the same level of detail that Goldstein does in 9.6, but it is still good reading.)
- The reading on the Hamilton-Jacobi equations and Action-Angle Variables is **Goldstein** Ch.10 sections 10.1-10.6, and 10.8. We will cover more examples of this material on problem set #6.

## Problem Set 5

On this problem set there are 5 problems involving canonical transformations, Poisson brackets, and conserved quantities. In the last problem you will apply the Hamilton-Jacobi method to a problem for which you already know the solution.

### 1. Canonical Transformations [12 points]

In this problem we get some practice with canonical transformations from  $(q, p)$  to  $(Q, P)$ . We will also look at generating functions  $F(q, p, Q, P, t)$ , following the notation in Goldstein for  $F_1(q, Q, t)$ ,  $F_2(q, P, t)$ ,  $F_3(p, Q, t)$ , and  $F_4(p, P, t)$ .

- [2 points] Determine two possible generating functions for  $Q_i = q_i$  and  $P_i = p_i$ .
- [2 points] Find a generating function  $F_1(q, Q, t)$  for:  $Q = p/t$  and  $P = -qt$ .
- [4 points] For which parameters  $k, \ell, m, n$  is there a generating function  $F_1(q, Q)$  for:  $Q = q^k p^\ell$  and  $P = q^m p^n$ ?
- [4 points] For a particle with charge  $q$  and mass  $m$  moving in an electromagnetic field the Hamiltonian is given by

$$H = \frac{1}{2m}(\vec{p} - q\vec{A})^2 + q\phi \quad (1)$$

where  $\vec{A} = \vec{A}(\vec{x}, t)$  and  $\phi = \phi(\vec{x}, t)$  are the vector and scalar potentials. Here  $\{x_i, p_j\}$  are canonical coordinates and momenta.

Under a *gauge transformation* of the electromagnetic field:

$$\vec{A} \rightarrow \vec{A}' = \vec{A} + \vec{\nabla}f(\vec{x}, t), \quad \phi \rightarrow \phi' = \phi - \frac{\partial f(\vec{x}, t)}{\partial t},$$

while  $\vec{p} - q\vec{A}$  is unchanged. Show that this is a canonical transformation for the coordinates and momenta of a charged particle, and determine a generating function  $F_2(\vec{x}, \vec{P}, t)$  for this transformation.

### 2. Harmonic Oscillator [7 points] (Related to Goldstein Ch.9 #24)

- [2 points] For constant  $a$  and canonical variables  $\{q, p\}$ , show that the transformation

$$Q = p + iaq, \quad P = \frac{p - iaq}{2ia}$$

is canonical by using the theorem that allows you to check this by using Poisson brackets.

- [5 points] With a suitable choice for  $a$ , obtain a new Hamiltonian for the linear harmonic oscillator problem  $K = K(Q, P)$ . Solve the equations of motion with  $K$  to find  $Q(t)$ ,  $P(t)$ , and then find  $q(t)$  and  $p(t)$ .

### 3. Poisson Brackets and Conserved Quantities [4 points]

A system of two degrees of freedom is described by the Hamiltonian

$$H = q_1 p_1 - q_2 p_2 + a q_1^2 + b q_2^2,$$

with constants  $a$  and  $b$ . Show that  $u_1 = (p_1 + a q_1)/q_2$  and  $u_2 = q_1 q_2$  are constants of the motion.

### 4. Angular Momentum and the Laplace-Runge-Lenz vector [13 points]

Consider the angular momentum  $\vec{L} = \vec{x} \times \vec{p}$  for canonical variables  $\{x_i, p_j\}$  in 3-dimensions. The components can be written as  $L_i = \epsilon_{ijk} x_j p_k$  with an implicit sum on the repeated indices  $j$  and  $k$ . Here  $\epsilon_{ijk}$  is the Levi-Civita tensor

$$\epsilon_{ijk} = \begin{cases} +1 & \text{if } ijk = 123 \text{ or a cyclic combination of this} \\ -1 & \text{if } ijk = 321 \text{ or a cyclic combination of this} \\ 0 & \text{otherwise} \end{cases}$$

Often  $\epsilon_{ijk}$  is handy when we are considering cross-products:  $\vec{c} = \vec{a} \times \vec{b}$  is equivalent to  $c_i = \epsilon_{ijk} a_j b_k$ . Some properties you may find useful are:  $\epsilon_{ijk} = \epsilon_{jki}$ ,  $\epsilon_{jik} = -\epsilon_{ijk}$ , and  $\sum_k \epsilon_{ijk} \epsilon_{lmk} = \delta_{il} \delta_{jm} - \delta_{im} \delta_{jl}$ .

- (a) [4 points] As warm up, calculate the Poisson brackets  $[x_i, L_j]$ ,  $[p_i, L_j]$ ,  $[L_i, L_j]$ , and  $[L_i, \vec{L}^2]$ .

Now consider two particles attracted to each other by a central potential  $V(r) = -k/r$ , where  $r = |\vec{r}|$  is the distance between them. Taking the origin at the CM, the Hamiltonian for this system is  $H = \vec{p}^2/(2\mu) - k/r$  where  $\mu$  is the reduced mass and the  $r_i$  and  $p_j$  are canonical variables. The angular momentum,  $\vec{L} = \vec{r} \times \vec{p}$ , is conserved so you may assume that  $[L_i, H] = 0$  (some of you may recall proving this in 8.223).

- (b) [7 points] Show that the Laplace-Runge-Lenz vector,  $\vec{A} = \vec{p} \times \vec{L} - \mu k \vec{r}/r$ , is conserved.

Recall that the conservation of  $\vec{L}$  implies that the motion of the particles in this central force take place in a plane that is perpendicular to  $\vec{L}$ . The set of  $H, \vec{L}, \vec{A}$  gives 7 constants of motion, but for two particles there are at most 6 constants from integrating the equations of motion. Furthermore, at least one constant must refer to an initial time, and none of  $H, \vec{L}, \vec{A}$  do so. Hence there must be at least two relations between these constants. It is easy to see that  $\vec{L} \cdot \vec{A} = 0$  provides one relation.

- (c) [2 points] Show that the other relation is  $\vec{A}^2 = \mu^2 k^2 + 2\mu H \vec{L}^2$ .

[Read Goldstein section 3.9 to see how  $\vec{A}$  can be used to very easily find the orbital equation  $r = r(\theta)$  for motion in the plane.]

**5. An Exponential Potential** [13 points]

A particle with mass  $m = 1/2$  is moving along the  $x$ -axis inside a potential  $V(x) = \exp(x)$ , so its Hamiltonian is  $H = p^2 + e^x$ . You may assume  $p > 0$ .

- (a) [6 points] Determine a generating function  $F_2(x, P)$  that yields a new Hamiltonian  $K = P^2$ . (Feel free to check your results with mathematica.)
- (b) [3 points] What are the transformation equations  $P = P(x, p)$  and  $Q = Q(x, p)$ ?
- (c) [4 points] Determine  $x(t)$  and  $p(t)$ .

Question [not for points]: How would your analysis change if  $p < 0$ ?

**6. Projectile with Hamilton-Jacobi** [11 points] (Goldstein Ch.10 #17)

Solve the problem of the motion of a point projectile of mass  $m$  in a vertical plane using the Hamilton-Jacobi method. Find both the equation of the trajectory and the dependence of the coordinates on time. Assume that the projectile is fired off at time  $t = 0$  from the origin with the velocity  $v_0$ , making an angle  $\theta$  with the horizontal.

MIT OpenCourseWare  
<http://ocw.mit.edu>

8.09 Classical Mechanics III  
Fall 2014

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.