

MIT OpenCourseWare  
<http://ocw.mit.edu>

8.323 Relativistic Quantum Field Theory I  
Spring 2008

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY  
Physics Department

**8.323: Relativistic Quantum Field Theory I**

Prof. Alan Guth

May 7, 2008

**PROBLEM SET 10**

**REFERENCES:** Peskin and Schroeder, Sec. 3.6, Secs. 4.1-4.6, in addition to lecture slides that will be posted.

**Problem 1: Majorana Fermions** (*15 points*)

Problem 3.4 of Peskin and Schroeder.

**Problem 2: Supersymmetry** (*15 points*)

Problem 3.5 of Peskin and Schroeder.

**Problem 3: P, C, and T for Scalar and Dirac theories** (*10 points*)

Problem 3.7 of Peskin and Schroeder.

**Problem 4: Decay of a Scalar Particle** (*10 points*)

Problem 4.2 of Peskin and Schroeder.

**Problem 5 (Extra Credit): The Linked Cluster Theorem** (*10 points extra credit*)

The cancellation of disconnected diagrams that we found when we calculated  $\langle \Omega | T \{ \phi(x_1) \phi(x_2) \dots \phi(x_n) \} | \Omega \rangle$  is actually part of a more general “linked cluster” theorem that is applicable to any system for which the concept of an expectation value makes sense. If one wishes to calculate the expectation value of an exponential function, or of some quantity times an exponential function, then the linked cluster theorem is useful.

To express this theorem, let  $A_1, A_2, \dots$  denote “random quantities,” i.e., quantities for which expectation values can be defined. They might be quantum-mechanical operators, like  $\phi(x_1), \phi(x_2), \dots$ , or they might be classical random variables, such as the position or momentum of particles in a statistical mechanics ensemble. We assume that the operations of addition and multiplication are well-defined, and that the expectation value function  $\langle \quad \rangle$  has the following two properties:

- (1)  $\langle \quad \rangle$  is linear:  $\langle A + \lambda B \rangle = \langle A \rangle + \lambda \langle B \rangle$  for any  $A, B$ .
- (2)  $\langle A_1 A_2 \dots \rangle$  is independent of the order of the factors.

Note that (2) is not usually valid for quantum-mechanical operators, but does hold for time-ordered products of operators. For time-ordered products it is the time arguments that control the order of evaluation, so the order in which the operators are written has no effect.

Our first task is to define the *connected part*, or *cumulant*, of a product of random quantities. For the product of two quantities, the definition will be

$$\langle AB \rangle_c = \langle AB \rangle - \langle A \rangle \langle B \rangle, \quad (5.1)$$

which is also called the correlator of  $A$  and  $B$ . If  $A$  and  $B$  are uncorrelated, then

$$\langle AB \rangle_c = 0. \quad (5.2)$$

A generalization of this concept can be defined by induction, but first we need to define what is meant by a partition of the integers  $\{1, 2, \dots, N\}$ . A partition of  $\{1, \dots, N\}$  into subsets is defined as a specific way of dividing the integers 1– $N$  into non-empty subsets. More formally, a partition of  $\{1, \dots, N\}$  is a list of sets  $S_1, S_2, \dots, S_k$ , with the properties

- i) Each  $S_\alpha$  is non-empty.
- ii) Each  $S_\alpha \subset \{1, \dots, N\}$ .
- iii)  $S_\alpha \cap S_\beta = \emptyset$  if  $\alpha \neq \beta$ .
- iv)  $\bigcup_{\alpha=1, \dots, k} S_\alpha = \{1, \dots, N\}$ .

Two partitions which differ only by the ordering of the subscript labels on the  $S$ 's are considered identical. Note that properties (iii) and (iv) above can be summarized by saying that each of the integers  $\{1, \dots, N\}$  belongs to one and only one set  $S_\alpha$ . Assuming that the connected part has been defined for all products of size less than  $N$ , the connected part for  $\langle A_1 \dots A_N \rangle$  is defined by

$$\begin{aligned} \langle A_1 \dots A_N \rangle_c &= \langle A_1 \dots A_N \rangle \\ &- \sum_{\text{all proper partitions of } \{1, \dots, N\}} \left\langle \prod_{i \in S_1} A_i \right\rangle_c \dots \left\langle \prod_{i \in S_k} A_i \right\rangle_c, \end{aligned} \quad (5.3)$$

where “proper partitions” excludes the case  $k = 1$ ,  $S_1 = \{1, \dots, N\}$ . The definition (5.3) applies even when some or all of the  $A_i$ 's are equal; for example, Eq. (5.1) implies that  $\langle A^2 \rangle_c = \langle A^2 \rangle - \langle A \rangle^2$ .

- (a) Use this definition to calculate  $\langle ABC \rangle_c$  in terms of ordinary expectation values. Find also the expression for  $\langle A^3 \rangle_c$  in terms of ordinary expectation values.
- (b) Generalize Eq. (5.2) by showing that if two sets of random quantities  $A_1, \dots, A_M$  and  $B_1, \dots, B_N$  are independent of each other in the sense that

$$\langle A_{i_1} \dots A_{i_m} B_{j_1} \dots B_{j_n} \rangle = \langle A_{i_1} \dots A_{i_m} \rangle \langle B_{j_1} \dots B_{j_n} \rangle \quad (5.4)$$

for any subset of the  $A$ 's and  $B$ 's, then

$$\langle A_{i_1} \dots A_{i_m} B_{j_1} \dots B_{j_n} \rangle_c = 0 \quad (5.5)$$

whenever  $m$  and  $n$  are both greater than or equal to one. [*Hint*: Mathematical induction is likely to be useful here.]

We now consider expectation values of products of random quantities and exponentials of random quantities. The quantities in the product can be labeled  $A_1, A_2, \dots, A_N$ , and the argument of the exponential can be written as

$$B \equiv \sum_{i=1}^M \lambda_i A_i, \quad (5.6)$$

where  $M \geq N$  and  $M$  could be infinite. (Since the  $A_i$ 's had no particular ordering before writing this expression, there is no loss in generality in labeling a particular subset of the  $A_i$  as  $A_1, \dots, A_N$ .) Our goal is to prove the identity

$$\begin{aligned} \langle A_1 \dots A_N e^B \rangle &= \exp \{ \langle e^B \rangle_c - 1 \} \\ &\times \sum_{\substack{\text{all partitions} \\ \text{of } \{1, \dots, N\} \\ \text{into subsets}}} \left\langle \left( \prod_{i \in S_1} A_i \right) e^B \right\rangle_c \dots \left\langle \left( \prod_{i \in S_k} A_i \right) e^B \right\rangle_c. \end{aligned} \quad (5.7)$$

We will prove this by a method that is somewhat indirect.

Instead of trying to directly prove Eq. (5.7) from the definition (5.3), we instead introduce a new definition of connected part to help us complete the proof. We use  $\bar{c}$  to denote the new definition of connected part, which is defined by a special case of Eq. (5.7):

$$\left\langle e^{\sum_i \lambda_i A_i} \right\rangle \equiv \exp \left\{ \left\langle e^{\sum_i \lambda_i A_i} \right\rangle_{\bar{c}} - 1 \right\}, \quad (5.8)$$

where

$$\left\langle e^{\sum_i \lambda_i A_i} \right\rangle_{\bar{c}} \quad (5.9)$$

is to be viewed as a shorthand for

$$\sum_{n=0}^{\infty} \frac{1}{n!} \sum_{i_1} \cdots \sum_{i_n} \lambda_{i_1} \cdots \lambda_{i_n} \langle A_{i_1} \cdots A_{i_n} \rangle_{\bar{c}}. \quad (5.10)$$

We define  $\langle A_{i_1} \cdots A_{i_n} \rangle_{\bar{c}}$  to be independent of the ordering of the factors, so any terms in the above sum that differ only by ordering are to be considered equivalent.

To use Eq. (5.8) as a definition, consider expanding both sides to some fixed order in the  $\lambda$ 's. To zero order in  $\lambda$ ,

$$\begin{aligned} 1 &= \exp\{\langle 1 \rangle_{\bar{c}} - 1\} \\ \implies \langle 1 \rangle_{\bar{c}} &= 1. \end{aligned} \quad (5.11)$$

To first order,

$$\begin{aligned} \sum_i \lambda_i \langle A_i \rangle &= \sum_i \lambda_i \langle A_i \rangle_{\bar{c}} \\ \implies \langle A_i \rangle_{\bar{c}} &= \langle A_i \rangle. \end{aligned} \quad (5.12)$$

- (c) Continue expanding Eq. (5.8) through third order in the  $\lambda_i$ 's, verifying that  $\langle AB \rangle_{\bar{c}} = \langle AB \rangle_c$  and that  $\langle ABC \rangle_{\bar{c}} = \langle ABC \rangle_c$ .
- (d) Now use a proof by induction to show that

$$\begin{aligned} \langle A_1 \cdots A_N e^B \rangle &= \exp\{\langle e^B \rangle_{\bar{c}} - 1\} \\ &\times \sum_{\substack{\text{all partitions} \\ \text{of } \{1, \dots, N\} \\ \text{into subsets}}} \left\langle \left( \prod_{i \in S_1} A_i \right) e^B \right\rangle_{\bar{c}} \cdots \left\langle \left( \prod_{i \in S_k} A_i \right) e^B \right\rangle_{\bar{c}}. \end{aligned} \quad (5.13)$$

Note that this formula differs from Eq. (5.7) only by using  $\bar{c}$  rather than  $c$  to define the connected part. The definition (5.8) guarantees that Eq. (5.13) holds for  $N = 0$ . You can therefore assume that Eq. (5.13) holds for some  $N$ , and show that it must hold for  $N + 1$ .

- (e) Having derived Eq. (5.13), now use the special case  $\lambda_i = 0$  for all  $i$  to show that

$$\langle A_1 \cdots A_N \rangle_{\bar{c}} = \langle A_1 \cdots A_N \rangle_c \quad (5.14)$$

for all  $N$ .

- (f) To see how this general result applies to a free scalar quantum field theory, define  $\langle X \rangle \equiv \langle 0 | T(X) | 0 \rangle$ , where  $T$  is the time-ordered product. Suppose that  $A = \phi^2(x_1)$  and  $B = \phi^2(x_2)$ . Show that

$$\begin{aligned}
 \langle AB \rangle &= \text{diagram 1} + \text{diagram 2} + \text{diagram 3} \\
 &= 2\Delta_F(x_1 - x_2)^2 + \Delta_F(0)^2 \quad , \tag{5.15}
 \end{aligned}$$

and that  $\langle AB \rangle_c$  is given by the first graph alone.

- (g) Generalize the result in (f) to show that if each  $A_i$  is a normal-ordered product of free fields at the point  $x_i$ , then  $\langle A_1 \dots A_N \rangle_c$  is given by the sum of all Wick contractions that correspond to connected diagrams. [*Hint*: Again, a proof by induction is appropriate. Assume that the statement holds for all values of  $N$  less than some  $N_0$ , and then use Eq. (5.3) to show that the only diagrams that make a net contribution to  $\langle A_1 \dots A_N \rangle_c$  are the connected ones.

Once we identify the connected part  $\langle A_1 \dots A_N \rangle_c$  with connected graphs, as you have done in part (g), note that one can take  $B = -i \int d^4z \mathfrak{H}_I(z)$ , and then Eq. (5.7) becomes a generalization of Peskin and Schroeder's Eq. (4.53):

$$\left\langle 0 \left| T \left\{ \phi_I(x) \phi_I(y) e^{-i \int d^4z \mathfrak{H}_I(z)} \right\} \right| 0 \right\rangle$$

is given by the sum of all connected graphs times the exponential of all disconnected graphs.