

Chapter 10

Strong P, T Problem

One result of the analysis is that a term $\approx \int tr G \tilde{G}$ in the action of QCD is not negligible, even though, it is a total derivative (so it does not appear in the classical equations of motion).

Since it is a dimension 4 interaction, it should appear in our canonical form for QCD. Unfortunately it violates P and T , so its processes with a substantial coefficient would not agree with experience. Detailed estimates suggest

$$\left| \frac{\theta}{2\pi} \right| \quad (\text{the natural parameter}) \leq 10^{-9} \quad (10.1)$$

Since there is P and T violation in the electroweak interaction, this term will be needed for renormalization so unless there is an extra symmetry to explain why $\theta \approx 0$ (or can be rotated away) we have an ugly situation.

The required symmetry (Peccei-Quinn) is essentially axial baryon number, since this counts (Δ winding) and thus powers of $e^{i\theta}$.

If there is a symmetry of this type and spontaneously broken at a high scale F , the kinetic term of its Nambu-Goldstone phase field reads

$$\partial_\mu \varphi \partial_\mu \varphi^+ \xrightarrow{\varphi = F e^{i\theta}} F^2 (\partial_\mu \theta)^2 \quad (10.2)$$

This puts us back at our earlier analysis. We get, for large F a very light, very weakly interacting particle, the axion.

It is a serious candidate to supply the dark matter of the universe.

For more on axions, see my '83 Erice lectures.