

# CHAPTER 1 INTRODUCTION TO CALCULUS

## 1.1 Velocity and Distance (page 6)

Starting from  $f(0) = 0$  at constant velocity  $v$ , the distance function is  $f(t) = vt$ . When  $f(t) = 55t$  the velocity is  $v = 55$ . When  $f(t) = 55t + 1000$  the velocity is still 55 and the starting value is  $f(0) = 1000$ . In each case  $v$  is the slope of the graph of  $f$ . When  $v(t)$  is negative, the graph of  $f(t)$  goes downward. In that case area in the  $v$ -graph counts as negative.

Forward motion from  $f(0) = 0$  to  $f(2) = 10$  has  $v = 5$ . Then backward motion to  $f(4) = 0$  has  $v = -5$ . The distance function is  $f(t) = 5t$  for  $0 \leq t \leq 2$  and then  $f(t)$  equals  $5(4-t)$  (not  $-5t$ ). The slopes are 5 and -5. The distance  $f(3) = 5$ . The area under the  $v$ -graph up to time 1.5 is 7.5. The domain of  $f$  is the time interval  $0 \leq t \leq 4$ , and the range is the distance interval  $0 \leq f \leq 10$ . The range of  $v(t)$  is only 5 and -5.

The value of  $f(t) = 3t + 1$  at  $t = 2$  is  $f(2) = 7$ . The value 19 equals  $f(6)$ . The difference  $f(4) - f(1) = 9$ . That is the change in distance, when 4 - 1 is the change in time. The ratio of those changes equals 3, which is the slope of the graph. The formula for  $f(t) + 2$  is  $3t + 3$  whereas  $f(t + 2)$  equals  $3t + 7$ . Those functions have the same slope as  $f$ : the graph of  $f(t) + 2$  is shifted up and  $f(t + 2)$  is shifted to the left. The formula for  $f(5t)$  is  $15t + 1$ . The formula for  $5f(t)$  is  $15t + 5$ . The slope has jumped from 3 to 15.

The set of inputs to a function is its domain. The set of outputs is its range. The functions  $f(t) = 7 + 3(t-2)$  and  $f(t) = vt + C$  are linear. Their graphs are straight lines with slopes equal to 3 and  $v$ . They are the same function, if  $v = 3$  and  $C = 1$ .

$$1 v = 30, 0, -30; v = -10, 20 \quad 3 v(t) = \begin{cases} 2 & \text{for } 0 < t < 10 \\ 1 & \text{for } 10 < t < 20 \\ -3 & \text{for } 20 < t < 30 \end{cases} \quad v(t) = \begin{cases} 0 & \text{for } 0 < t < T \\ \frac{1}{T} & \text{for } T < t < 2T \\ 0 & \text{for } 2T < t < 3T \end{cases}$$

$$5 25; 22; t+10 \quad 7 6; -30 \quad 9 v(t) = \begin{cases} 20 & \text{for } t < .2 \\ 0 & \text{for } t > .2 \end{cases} \quad f(t) = \begin{cases} 20t & \text{for } t \leq .2 \\ 4 & \text{for } t \geq .2 \end{cases} \quad 11 10\%; 12\frac{1}{2}\%$$

$$13 f(t) = 0, 30(t-1), 30; f(t) = -30t, -60, 30(t-6) \quad 15 \text{ Average } 8, 20 \quad 17 40t - 80 \text{ for } 1 \leq t \leq 2.5$$

$$21 0 \leq t \leq 3, -40 \leq f \leq 20; 0 \leq t \leq 3T, 0 \leq f \leq 60T \quad 23 3 - 7t \quad 25 6t - 2 \quad 27 3t + 7$$

$$29 \text{ Slope } -2; 1 \leq f \leq 9 \quad 31 v(t) = \begin{cases} 8 & \text{for } 0 < t < T \\ -2 & \text{for } T < t < 5T \end{cases} \quad f(t) = \begin{cases} 8t & \text{for } 0 \leq t \leq T \\ 10T - 2t & \text{for } T \leq t \leq 5T \end{cases}$$

$$33 \frac{9}{5}C + 32; \text{ slope } \frac{9}{5} \quad 35 f(w) = \frac{w}{1000}; \text{ slope} = \text{conversion factor} \quad 37 1 \leq t \leq 5, 0 \leq f \leq 2$$

$$39 0 \leq t \leq 5, 0 \leq f \leq 4 \quad 41 0 \leq t \leq 5, 1 \leq t \leq 32 \quad 43 \frac{1}{2}t + 4; \frac{1}{2}t + \frac{7}{2}; 2t + 12; 2t + 3$$

$$45 \text{ Domains } -1 \leq t \leq 1 : \text{ ranges } 0 \leq 2t + 2 \leq 4, \quad -3 \leq t - 2 \leq -1, \quad -2 \leq -f(t) \leq 0, \quad 0 \leq f(-t) \leq 2$$

$$47 \frac{3}{2}V; \frac{3}{2}V \quad 49 \text{ input * input} \rightarrow A \quad \text{input * input} \rightarrow A \quad B * B \rightarrow C \quad \text{input} + 1 \rightarrow A \\ \text{input} + A \rightarrow \text{output} \quad \text{input} + A \rightarrow B \quad B + C \rightarrow \text{output} \quad A * A \rightarrow B \\ A + B \rightarrow \text{output}$$

$$51 3t + 5, 3t + 1, 6t - 2, 6t - 1, -3t - 1, 9t - 4; \text{ slopes } 3, 3, 6, 6, -3, 9$$

$$53 \text{ The graph goes up and down twice. } f(f(t)) = \begin{cases} 2(2t) & 0 \leq t \leq 1.5 \\ 12 - 4t & 1.5 \leq t \leq 3 \end{cases} \quad \begin{cases} 12 - 2(12 - 2t) & 3 \leq t \leq 4.5 \\ 2(12 - 2t) & 4.5 \leq t \leq 6 \end{cases}$$

2 (a) The slopes are  $v = 2$  then  $v = 1$  then  $v = -3$

(b) The slopes are  $v = 0$  then  $v = 1/T$  then  $v = 0$

**4**  $f(t) = 20(t - 1)$  for  $1 \leq t \leq 2$

**6**  $f(1.4T) = .4$ ; if  $T = 3$  then  $f(4) = \frac{1}{3}$ . This is  $\frac{1}{3}$  of the distance between  $f(3) = 0$  and  $f(6) = 1$ .

**8** Average speed =  $\frac{f(2)-f(0)}{2} = \frac{20-10}{2} = 5$ ; the average speed is zero between  $t = \frac{1}{2}$  and  $t = 1\frac{1}{4}$ , since at both times  $f = 5$ .

**10**  $v(t)$  is negative-zero-positive;  $v(t)$  is above 55 then equal to 55;  $v(t)$  increases in jumps;  $v(t)$  is zero then positive. All with corresponding  $f(t)$ .

**12**  $f(t)$  increases linearly from 5.2 billion in 1990 to 6.2 billion in 2000.

**14** (a)  $f(t) = -40t$  (graph drops linearly to -40 at  $t = 1$ ) then  $f(t) = -40 + 40(t - 1) = 40t - 80$ .

End at  $f(\frac{5}{2}) = 20$

(b) Second graph rises to  $40T$  at time  $T$ , stays constant until time  $2T$ , then rises more slowly to  $60T$  at time  $3T$ .

**16**  $f(t) = \begin{cases} 0 & 0 \leq t \leq 1 \\ 30(t-1) & 1 \leq t \leq 2; \\ 30 & t \geq 2 \end{cases} f(t) = \begin{cases} -30t & 0 \leq t \leq 2 \\ -60 & 2 \leq t \leq 4 \\ -60 + 30(t-4) & t > 4 \end{cases}$

**18**  $v(t) = 8$  then 1 (after  $t = 2$ );  $f(t) = 6 + 8t$  then  $20 + t$ .

**20**  $1200 + 30x = 40x$  when  $1200 = 10x$  or  $x = 120$  yearbooks. The slope is 30. If it goes above 40 you can't break even.

**22** Range = {0, 20, 40}; the velocity is not defined at the jump.

**24**  $f(t) = 4t + 1$  (linear up) or  $-4t + 9$  (linear down).

**26** The function increases by 2 in one time unit so the slope (velocity) is 2;  $f(t) = 2t + C$  with constant  $C = f(0)$ .

**28**  $f(2t) = 2vt$  must equal  $4vt$  so  $v = 0$  and  $f = 0$ . But  $\frac{1}{2}a(2t)^2$  does equal  $4(\frac{1}{2}at^2)$ . To go four times as far in twice the time, you must accelerate.

**30**  $f(t) = 0$  then  $8 - 2t$  (change at  $t = 4$ ); slopes 0 and -2; range  $-2 \leq f(t) \leq 0$ .

**32**  $f(t) = 3t = 12$  at  $t = 4$ ; then  $v = 6$  gives  $f(t) = 12 + 6(t-4) = 30$  at  $t = 7$ . The extra distance was 18 in 3 time units; thus  $v(t) = 3$  then 6.

**34**  $C(F) = \frac{5}{9}(F - 32)$  has slope  $\frac{5}{9}$ .

**36** At  $t = 0$  the reading was  $.061 + 10(.015) = .211$ . A drop of  $.061 - .04 = .021$  would take  $.021/.015$  hours. This was the Exxon Valdez accident.

**38** Domain  $1 < t \leq 5$ ; range  $\frac{1}{4} \leq f(t) < \infty$ .

**40** Domain  $0 \leq t < 4$  and  $4 < t \leq 5$  (omit  $t = 4$ ); range  $\frac{1}{16} \leq f(t) < \infty$

**42** Domain  $0 \leq t \leq 5$ ; range  $2^{-5}$  (or  $\frac{1}{32}$ )  $\leq f(t) \leq 1$ .

**44** Jump from 0 to 1 at  $t = 0$ ; jump from 2 to 3 at  $t = 0$ ; jump from 0 to 1 at  $t = -2$ ; jump from 0 to 3 at  $t = 0$ ; jump from 0 to 1 at  $t = 0$ .

**46**  $2f(3t) = 2(3t - 1) = 6t - 2$ ;  $f(1-t) = (1-t) - 1 = -t$ ;  $f(t-1) = (t-1) - 1 = t - 2$ .

**48**  $f_1(t) = 3t + 3$ ;  $f_2(t) = 3t + 18$ .

**50** "A function assigns an output to each input ..."

**52**  $3(vt + C) + 1$  has slope  $3v$ ;  $v(3t + 1) + C$  also has slope  $3v$ ;  $2(4vt + C)$  has slope  $8v$ ;  $-vt + C$  has slope  $-v$ ;  $vt + C - C$  has slope  $v$ ;  $v(vt + C) + C$  has slope  $v^2$ .

**54** A function cannot have two values (the upper and lower branches of X) at the same point. Apparently only

**U, V, W** are graphs. Their slopes are negative-positive and negative-positive-negative-positive.

## 1.2 Jumps in Velocity (page 14)

When the velocity jumps from  $v_1$  to  $v_2$ , the function  $v(t)$  is piecewise constant. The distance function  $f(t)$  is piecewise linear. In the first time interval,  $f(t) = f(0) + v_1 t$ . After the jump at  $t = 1$ , the formula is  $f(t) = f(1) + v_2(t - 1)$ . In case  $f_0 = 6$  all distances are increased by 6 and all velocities are the same.

With distances 1, 5, 25 at unit times, the velocities are **4** and **20**. These are the slopes of the  $f$ -graph. The slope of the tax graph is the tax rate. If  $f(t)$  is the postage cost for  $t$  ounces or  $t$  grams, the slope is the cost per ounce (or per gram). For distances 0, 1, 4, 9 the velocities are 1, 3, 5. The sum of the first  $j$  odd numbers is  $f_j = j^2$ . Then  $f_{10}$  is 100 and the velocity  $v_{10}$  is 19.

The piecewise linear sine has slopes 1, 0, -1, -1, 0, 1. Those form a piecewise constant cosine. Both functions have period equal to 6, which means that  $f(t + 6) = f(t)$  for every  $t$ . The velocities  $v = 1, 2, 4, 8, \dots$  have  $v_j = 2^{j-1}$ . In that case  $f_0 = 1$  and  $f_j = 2^j$ . The sum of 1, 2, 4, 8, 16 is 31. The difference  $2^j - 2^{j-1}$  equals  $2^{j-1}$ . After a burst of speed  $V$  to time  $T$ , the distance is  $VT$ . If  $f(T) = 1$  and  $V$  increases, the burst lasts only to  $T = 1/V$ . When  $V$  approaches infinity,  $f(t)$  approaches a step function. The velocities approach a delta function, which is concentrated at  $t = 0$  but has area 1 under its graph. The slope of a step function is zero or infinity.

- 1** 1.1, -2, 5      **3** 6.6, 8.8; -11, -15; 4, 14      **5**  $h(t) = 9t + 6$ , add slopes      **7**  $f = 2t$  then  $3t - T$   
**9** 7, 28,  $8t + 4$ ; multiply slopes      **11** 16, 0,  $8t$  then  $36 - 4t$       **13** Tax =  $.28x$ ; 280,000      **15**  $19\frac{1}{4}\%$   
**17** All  $v_j = 2$ ;  $v_j = (-1)^{j-1}$ ;  $v_j = (\frac{1}{2})^j$       **21**  $j^2 + j$       **23**  $f_{10} = 38$       **25**  $(101^2 - 99^2)/2 = \frac{400}{2}$   
**27**  $v_j = 2j$       **29**  $f_{31} = 5$       **31**  $a_j = -f_j$       **33** 0; 1; .1      **35** require  $v_2 = -v_1$   
**37**  $v_j = 3(4)^{j-1}$       **39**  $v_j = -(\frac{1}{2})^j$       **41**  $v_j = 2(-1)^j$ , sum is  $f_j - 1$       **45**  $v = 1000, t = 10/V$   
**47** M, N      **51**  $\sqrt{9} < 2 \cdot 9 < 9^2 < 2^9$ ;  $(\frac{1}{9})^2 < 2(\frac{1}{9}) < \sqrt{1/9} < 2^{1/9}$

**2**  $f(6), f(7)$  are 66, 77 and -11, -13 and 4, 9. Then  $f(7) - f(6)$  is **11, -2, 5**.

**4** The increases  $f(4) - f(1)$  are  $12 - 3 = 9$  and  $14 - 5 = 9$  and  $18 - 9 = 9$ .

**6**  $h(t) = .5t + 3$ ; the slopes of  $f, g, h$  are **3, 2.5** and  $3 - 2.5 = .5$ .

**8**  $f(t) = 1 + 10t$  for  $0 \leq t \leq \frac{1}{10}$ ,  $f(t) = 2$  for  $t \geq \frac{1}{10}$

**10**  $f(3) = 12$ ;  $g(f(3)) = g(12) = 25$ ;  $g(f(t)) = g(4t) = 8t + 1$ . Distance increases four times as fast and velocity is multiplied by **4**.

**12** 10,160.50 is  $f(44,900) = 2782.50 + .28(44,900 - 18,550)$ .

**14**  $F(x) = 2f(\frac{1}{2}x) = .15x$  for  $x \leq 37,100$ ; then  $F(x) = 5565 + .28(x - 37,100)$  up to  $x = 89,800$ ; then  $F(x) = 20,321 + .33(x - 89,800)$  up to  $x = 186,260$ ; then  $F(x) = .28x$  beyond 186,260.

The 1991 rates on the front cover have only three brackets.

**16**  $f(t) = 3 + 2t$  for  $t \geq 1$  is continuous;  $f(t) = 4 + 2t$  is discontinuous (because  $f(1) = 5$ ).  $f(x) = .15x$

- then  $3000 + .28(x - 18,550)$  has a jump at \$18,550.
- 18  $f_1 = 1, f_2 = 3, f_3 = 7, f_j = 2^j - 1; f_1 = -1, f_2 = 0, f_3 = -1, f_j = \{-1 \text{ for odd } j, 0 \text{ for even } j\}$   
 $= \frac{1}{2}((-1)^j - 1).$
- 20 The big triangle has area  $= \frac{1}{2}(\text{base})(\text{height}) = \frac{1}{2}j^2$  and the  $j$  small triangles have area  $\frac{1}{2}j$ . Together they give rectangles of total area  $1 + 2 + \dots + j$ . Note: Another drawing could move the diagonal line up by  $\frac{1}{2}$ . The big triangle still has area  $\frac{1}{2}j^2$  and the strip across the bottom has area  $\frac{1}{2}j$ .
- 22 False when the  $v_j$  are  $(\frac{1}{2})^j$ ; false when the  $v_j$  are  $-(\frac{1}{2})^j$ ; true when all  $f_{j+p} = f_j$  ( $p$  is the period) because then  $v_{j+p} = f_{j+p} - f_{j+p-1}$  equals  $f_j - f_{j-1} = v_j$ ; false when all  $v_j = 1$ .
- 24 Assume  $f_0 = 0$ . First  $f_j = j^2$ , second  $f_j = j$ , by addition third  $f_j = j^2 + j$ , by division last  $f_j = \frac{1}{2}(j^2 + j)$  which is  $1 + 2 + \dots + j$ .
- 26  $f(99) = 9900$  and  $f(101) = 10302; \Delta f/\Delta t = 402/2 = 201$ .
- 28 Take  $v = C, 2C, 3C, \dots$  Then  $f = C, 3C, 6C, \dots$ . The example  $f_3 - 2f_2 + f_1$  gives  $6C - 2(3C) + C = C$ . The answer is always  $C$  (by Problem 30).
- 30  $f_{j+1} - 2f_j + f_{j-1}$  equals  $(f_{j+1} - f_j) - (f_j - f_{j-1}) = v_{j+1} - v_j$ . If  $v$  is velocity then  $a$  is acceleration.
- 32 The period of  $v + w$  is 30, the smallest multiple of both 6 and 10. (Then  $v$  completes five cycles and  $w$  completes three.) An example for functions is  $v = \sin \frac{\pi x}{3}$  and  $w = \sin \frac{\pi x}{5}$  ( $v + w$  has a nice graph).
- 34  $f(12) = (1 + 2 + 1 + 0) + (1 + 2 + 1 + 0) + (1 + 2 + 1 + 0) = 12$ . Then  $f(14) = 12 + 1 + 2 = 15$  and  $f(16) = 15 + 1 + 0 = 16$ .  $f$  doesn't have period 4 since  $x_1 + x_2 + x_3 + x_4$  is not zero.
- 36  $2^j$  is 2 times  $2^{j-1}$ . Subtracting  $2^{j-1}$  leaves  $2^{j-1}$ . Similarly  $3^j$  is 3 times  $3^{j-1}$  and subtraction leaves 2 times  $3^{j-1}$ .
- 38  $f_1 - f_0$  equals  $v_1 = 2f_0 = 2$  so  $f_1$  is 3;  $f_2 - f_1$  equals  $v_2 = 2f_1 = 6$  so  $f_2$  is 9; then  $f_3$  is 27 and  $f_4$  is 81. Problem 36 shows that  $f_j = 3^j$  fits the requirement  $v_j = 2f_{j-1}$ .
- 40  $v_j = f_j - f_{j-1}$  equals  $r^j - r^{j-1}$ . Adding the  $v$ 's gives  $(f_1 - f_0) + (f_2 - f_1) + (f_3 - f_2) + \dots + (f_j - f_{j-1})$ . Cancelling leaves only  $f_j - f_0 = r^j - 1$ .
- 42 The first sum is  $1024 - 1 = 1023$ . The second is  $2 - \frac{1}{512} = \frac{1023}{512}$ . (Notice how the second sum is  $\frac{1}{512}$  times the first.) The sum formula is in Problem 43 and also Problem 18.
- 44  $U(t) - U(t - 1)$  is zero except between  $t = 0$  and  $t = 1$  (where it equals 1). If this is the velocity, then the distance is  $f(t) = t$  up to  $t = 1$ ; then  $f(t) = 1$ : a "short burst of speed". If the square wave is distance, then  $v(t)$  is a delta function at  $t = 0$  minus a delta function at  $t = 1$ .
- 46 The sum jumps up by 1 at  $t = 0, 1, 2$ . Its slope is a sum of three delta functions.
- 48  $\begin{cases} \text{For } j = 1, N \text{ do} \\ v_j = f_j - f_{j-1} \end{cases}$  Examples  $2^j$  and  $j^2$  and  $2^j$  give  $v_j = 2$  and  $v_j = 2j - 1$  and  $v_j = 2^{j-1}$ .
- 50 FINDV (FINDF ( $v_1, \dots, v_N$ )) brings back  $v_1, \dots, v_N$ . But FINDF (FINDV ( $f_0, f_1, \dots, f_N$ )) produces  $0, f_1 - f_0, f_2 - f_0, \dots, f_N - f_0$ .
- 52 The average age increases with slope 1 except at a birth or death (when it is discontinuous).

### 1.3 The Velocity at an Instant (page 21)

Between the distances  $f(2) = 100$  and  $f(6) = 200$ , the average velocity is 25. If  $f(t) = \frac{1}{4}t^2$  then  $f(6) = 9$  and  $f(8) = 16$ . The average velocity in between is 3.5. The instantaneous velocities at  $t = 6$  and  $t = 8$  are 3

and 4.

The average velocity is computed from  $f(t)$  and  $f(t + h)$  by  $v_{ave} = \frac{1}{h}(f(t + h) - f(t))$ . If  $f(t) = t^2$  then  $v_{ave} = 2t + h$ . From  $t = 1$  to  $t = 1.1$  the average is 2.1. The instantaneous velocity is the limit of  $v_{ave}$ . If the distance is  $f(t) = \frac{1}{2}at^2$  then the velocity is  $v(t) = at$  and the acceleration is  $a$ .

On the graph of  $f(t)$ , the average velocity between  $A$  and  $B$  is the slope of the secant line. The velocity at  $A$  is found by letting  $B$  approach  $A$ . The velocity at  $B$  is found by letting  $A$  approach  $B$ . When the velocity is positive, the distance is increasing. When the velocity is increasing, the car is accelerating.

- 1** 6, 6,  $\frac{13}{2}a$ ,  $-12$ , 0, 13    **3** 4, 3.1,  $3 + h$ , 2.9    **5** Velocity at  $t = 1$  is 3    **7** Area  $f = t + t^2$ , slope of  $f$  is  $1 + 2t$   
**9** F; F; F; T    **11** 2;  $2t$     **13**  $12 + 10t^2$ ;  $2 + 10t^2$     **15** Time 2, height 1, stays above  $\frac{3}{4}$  from  $t = \frac{1}{2}$  to  $\frac{3}{2}$   
**17**  $f(6) = 18$     **21**  $v(t) = -2t$  then  $2t$     **23** Average to  $t = 5$  is 2;  $v(5) = 7$     **25**  $4v(4t)$     **27**  $v_{ave} = t$ ,  $v(t) = 2t$

- 2** (a)  $\frac{6(t+h)-6t}{h} = 6$  (limit is 6); (b)  $\frac{6(t+h)+2-(6t+2)}{h} = 6$  (limit also 6); (c)  $\frac{\frac{1}{2}a(t^2+2th+h^2)-\frac{1}{2}at^2}{h} = at + \frac{1}{2}ah$  (limit is  $at$ ); (d)  $\frac{t+h-(t+h)^2-(t-t)^2}{h} = 1 - 2t - h$  (limit is  $1 - 2t$ ); (e)  $\frac{6-6}{h} = 0$  (limit is 0); (f) the limit is  $v(t) = 2t$  (and  $f(t) = t^2$  gives  $\frac{(t+h)^2-t^2}{h} = 2t + h$ ).  
**4**  $\frac{\Delta f}{\Delta t} = \frac{2-0}{1} = 2$ ;  $\frac{3/4-0}{1/2} = \frac{3}{2}$ ;  $\frac{h+h^2-0}{h} = 1 + h$ .    **6**  $\lim \frac{\Delta f}{\Delta t} = \lim(1 + h) = 1 =$  slope of the parabola at  $t = 0$ .  
**8**  $v(t) = 3 - 2t$  gives a line through (0,3) and (1,1);  $f(t) = 3t - t^2$  gives a parabola through (0,0) and (3,0) with maximum at  $(\frac{3}{2}, \frac{9}{4})$ .  
**10** Slope of  $f(t) = 6t^2$  is  $v(t) = 12t$ ; slope of  $v(t) = 12t$  is  $a = 12 =$  acceleration.  
**12**  $\Delta f = \frac{1}{2}a(t+h)^2 - \frac{1}{2}a(t-h)^2 = 2ath$ ; then  $\frac{\Delta f}{\Delta t} = \frac{2ath}{2h} = at =$  velocity at time  $t$ . The region under the line  $v = at$  is a trapezoid. Its area is the base  $2h$  times the average height  $at$ .  
**14** True (the slope is  $\frac{\Delta f}{\Delta t}$ ); false (the curve is partly steeper and partly flatter than the secant line which gives the average slope); true (because  $\Delta f = \Delta F$ ); false ( $V$  could be larger than  $v$  in between).  
**16** The functions are  $t^2$  and  $t^2 - 2$  and  $4t^2$ . The velocities are  $2t$  and  $2t$  and  $8t$ .  
**18** The graph is a parabola  $f(t) = \frac{1}{2}t^2$  out to  $f = 2$  at  $t = 2$ . After that the slope of  $f$  stays constant at 2.  
**20** Area to  $t = 1$  is  $\frac{1}{2}$ ; to  $t = 2$  is  $\frac{3}{2}$ ; to  $t = 3$  is 2; to  $t = 4$  is  $\frac{3}{2}$ ; to  $t = 5$  is  $\frac{1}{2}$ ; area from  $t = 0$  to  $t = 6$  is zero. The graph of  $f(t)$  through these points is parabola-line-parabola (symmetric)-line-parabola to zero.  
**22**  $f(t)$  is a parabola  $t - \frac{1}{2}t^2$  through (0,0),  $(1, \frac{1}{2})$ , and  $(2,0)$ ;  $f(t)$  is the same parabola until  $(1, \frac{1}{2})$ , but the second half goes up to  $(2,1)$ ;  $f(t)$  is the parabola  $2t - t^2$  until  $(1,1)$  and then a horizontal line since  $v = 0$ .  
**24** The slope of  $f$  is  $v(t) = at + b$ ; the slope of  $v$  is the constant  $a$ ;  $f(t) = \frac{1}{2}t^2 + t + 1$  equals 41 when  $t = 8$ . (The quadratic formula for  $\frac{1}{2}t^2 + t - 40 = 0$  gives  $t = -1 \pm \sqrt{1^2 + 80} = -1 \pm 9$ .)  
**26**  $f(t) = t - t^2$  has  $v(t) = 1 - 2t$  and  $f(3t) = 3t - 9t^2$ . The slope of  $f(3t)$  is  $3 - 18t$ . This is  $3v(3t)$ .  
**28** To find  $f(t)$  multiply the time  $t$  by the average velocity. This is because  $v_{ave}(t) = \frac{f(t)-f(0)}{t} = \frac{f(t)}{t}$ .

## 1.4 Circular Motion (page 28)

A ball at angle  $t$  on the unit circle has coordinates  $x = \cos t$  and  $y = \sin t$ . It completes a full circle at  $t = 2\pi$ . Its speed is 1. Its velocity points in the direction of the tangent, which is perpendicular to the radius

coming out from the center. The upward velocity is  $\cos t$  and the horizontal velocity is  $-\sin t$ .

A mass going up and down level with the ball has height  $f(t) = \sin t$ . This is called simple harmonic motion. The velocity is  $v(t) = \cos t$ . When  $t = \pi/2$  the height is  $f = 1$  and the velocity is  $v = 0$ . If a speeded-up mass reaches  $f = \sin 2t$  at time  $t$ , its velocity is  $v = 2 \cos 2t$ . A shadow traveling *under* the ball has  $f = \cos t$  and  $v = -\sin t$ . When  $f$  is distance = area = integral,  $v$  is velocity = slope = derivative.

- 1**  $10\pi, (0, -1), (-1, 0)$       **3**  $(4 \cos t, 4 \sin t); 4$  and  $4t; 4 \cos t$  and  $-4 \sin t$   
**5**  $3t; (\cos 3t, \sin 3t); -3 \sin 3t$  and  $3 \cos 3t$       **7**  $x = \cos t; \sqrt{2}/2; -\sqrt{2}/2$       **9**  $2\pi/3; 1; 2\pi$   
**11** Clockwise starting at  $(1, 0)$       **13** Speed  $\frac{2}{\pi}$       **15** Area 2      **17** Area 0  
**19** 4 from speed, 4 from angle      **21**  $\frac{1}{4}$  from radius times 4 from angle gives 1 in velocity  
**23** Slope  $\frac{1}{2}$ ; average  $(1 - \frac{\sqrt{3}}{2})/(\pi/6) = \frac{3(2-\sqrt{3})}{\pi} = .256$       **25** Clockwise with radius 1 from  $(1, 0)$ , speed 3  
**27** Clockwise with radius 5 from  $(0, 5)$ , speed 10      **29** Counterclockwise with radius 1 from  $(\cos 1, \sin 1)$ , speed 1  
**31** Left and right from  $(1, 0)$  to  $(-1, 0)$ ,  $v = -\sin t$       **33** Up and down between 2 and  $-2$ ; start  $2 \sin \theta$ ,  $v = 2 \cos(t+\theta)$   
**35** Up and down from  $(0, -2)$  to  $(0, 2)$ ;  $v = \sin \frac{1}{2}t$       **37**  $x = \cos \frac{2\pi t}{360}, y = \sin \frac{2\pi t}{360}$ , speed  $\frac{2\pi}{360}$ ,  $v_{\text{up}} = \cos \frac{2\pi t}{360}$   
**39** I think there is a stop between backward and forward motion.

- 2** The cosine of  $\frac{2\pi}{3}$  is  $x = -\frac{1}{2}$ ; the sine is  $y = \frac{\sqrt{3}}{2}$ ; the tangent is  $\frac{y}{x} = -\sqrt{3}$ ; the ball has a distance  $\sqrt{3}$  to go (draw triangle from  $(0,0)$  to  $(x,y)$  and back down at right angle); the speed is 1 so the added time is  $\sqrt{3}$  and the total time is  $\frac{2\pi}{3} + \sqrt{3}$ . Not easy.  
**4**  $x = R \cos t$  and  $y = R \sin t$ ; velocity  $-R \sin t$  and  $R \cos t$ ; distance and velocity triangles both grow by  $R$ .  
**6** The angle is  $\frac{\pi}{2} + 3t$ ; the position is  $x = \cos(\frac{\pi}{2} + 3t) = -\sin 3t$  and  $y = \sin(\frac{\pi}{2} + 3t) = \cos 3t$ ; the vertical velocity is  $-3 \sin 3t$  (= horizontal velocity of original ball).  
**8** The new mass at  $x = \cos t, y = 0$  never meets the old mass at  $x = 0, y = \sin t$ . The distance between them is always  $\sqrt{\cos^2 t + \sin^2 t} = 1$ .  
**10**  $f = \sin(t + \pi)$  equals  $-\sin t$ ; the velocity is  $\cos(t + \pi)$  which equals  $-\cos t$ . The ball is a half-circle ahead of the original ball.  
**12**  $f(t) = \sin t + \cos t$  has  $f^2 = \sin^2 t + 2 \sin t \cos t + \cos^2 t$  which is the same as  $1 + 2 \sin t \cos t$  (or  $1 + \sin 2t$ ). The maximum is at  $t = 45^\circ = \frac{\pi}{4}$  when  $f^2 = 2$ . Then  $f_{\max} = \sqrt{2}$ . Its graph is a sine curve with this maximum point:  $f(t)$  equals  $\sqrt{2} \sin(t + \frac{\pi}{4})$ .  
**14** The ball goes halfway around the circle in time  $\pi$ . For the mass to fall a distance 2 in time  $\pi$  we need  $2 = \frac{1}{2}a\pi^2$  so  $a = 4/\pi^2$ .  
**16** The area is  $f(t) = \sin t$ , and  $\sin \frac{\pi}{6} - \sin 0 = \frac{1}{2}$ .  
**18** The area is still  $f(t) = \sin t$ , and  $\sin \frac{3\pi}{2} - \sin \frac{\pi}{2} = -1 - 1 = -2$ .  
**20** The radius is 2 and time is speeded up by 3 so the velocity is 6 with minus sign because the cosine starts downward (ball moving to left).  
**22** The distance is  $-\cos 5t$ .  
**24**  $\frac{\sin 1 - \sin 0}{1} = .8415$  and  $\frac{\sin .1}{.1} = .9983$  and  $\frac{\sin .01}{.01} = .9999$ ; then  $\frac{\sin .001}{.001} = .99999983$ .  
**26** Counterclockwise with radius 3 starting at  $(3, 0)$  with speed 12.  
**28** Counterclockwise with radius 1 around center at  $(1, 0)$ . Starts from  $(2, 0)$ ; speed 1.  
**30** Clockwise around the unit circle from  $(1, 0)$  with speed 1.  
**32** Up and down between  $-1$  and  $1$ , starting at  $(0, 0)$  with velocity  $5 \cos 5t$ .  
**34** Along the  $45^\circ$  line  $y = x$  between  $(-1, -1)$  and  $(1, 1)$ . Starting at  $(1, 1)$  with  $x$  and  $y$  velocities  $-\sin t$ .

**36** Along the line  $x + y = 1$  between  $(1,0)$  and  $(0,1)$ . Starting at  $(1,0)$  the  $x$  and  $y$  velocities are  $-2 \sin t \cos t$  and  $2 \sin t \cos t$ . (Maybe introduce  $\cos^2 t = \frac{1}{2} + \frac{1}{2} \cos 2t$  and  $\sin^2 t = \frac{1}{2} - \frac{1}{2} \cos 2t$  to find velocities  $-\sin 2t$  and  $\sin 2t$ : Discuss.)

**38** Choose  $k = 2\pi$ . The speed is  $2\pi$  and the upward velocity is  $2\pi \cos 2\pi t$ .

## 1.5 Review of Trigonometry (page 33)

Starting with a right triangle, the six basic functions are the ratios of the sides. Two ratios (the cosine  $x/r$  and the sine  $y/r$ ) are below 1. Two ratios (the secant  $r/x$  and the cosecant  $r/y$ ) are above 1. Two ratios (the tangent and the cotangent) can take any value. The six functions are defined for all angles  $\theta$ , by changing from a triangle to a circle.

The angle  $\theta$  is measured in radians. A full circle is  $\theta = 2\pi$ , when the distance around is  $2\pi r$ . The distance to angle  $\theta$  is  $\theta r$ . All six functions have period  $2\pi$ . Going clockwise changes the sign of  $\theta$  and  $\sin \theta$  and  $\tan \theta$ . Since  $\cos(-\theta) = \cos \theta$ , the cosine is unchanged (or even).

Coming from  $x^2 + y^2 = r^2$  are the three identities  $\sin^2 \theta + \cos^2 \theta = 1$  and  $\tan^2 \theta + 1 = \sec^2 \theta$  and  $1 + \cot^2 \theta = \csc^2 \theta$ . (Divide by  $r^2$  and  $x^2$  and  $y^2$ .) The distance from  $(2,5)$  to  $(3,4)$  is  $d = \sqrt{2}$ . The distance from  $(1,0)$  to  $(\cos(s-t), \sin(s-t))$  leads to the addition formula  $\cos(s-t) = \cos s \cos t + \sin s \sin t$ . Changing the sign of  $t$  gives  $\cos(s+t) = \cos s \cos t - \sin s \sin t$ . Choosing  $s = t$  gives  $\cos 2t = \cos^2 t - \sin^2 t$  or  $2 \cos^2 t - 1$ . Therefore  $\frac{1}{2}(1 + \cos 2t) = \cos^2 t$ , a formula needed in calculus.

**1** Connect corner to midpoint of opposite side, producing  $30^\circ$  angle      **3**  $\pi$       **7**  $\frac{\theta}{2\pi} \rightarrow$  area  $\frac{1}{2}r^2\theta$

**9**  $d = 1$ , distance around hexagon < distance around circle      **11** T; T; F; F

**13**  $\cos(2t+t) = \cos 2t \cos t - \sin 2t \sin t = 4 \cos^3 t - 3 \cos t$

**15**  $\frac{1}{2}\cos(s-t) + \frac{1}{2}\cos(s+t); \frac{1}{2}\cos(s-t) - \frac{1}{2}\cos(s+t)$       **17**  $\cos \theta = \sec \theta = \pm 1$  at  $\theta = n\pi$

**19** Use  $\cos(\frac{\pi}{2} - s - t) = \cos(\frac{\pi}{2} - s) \cos t + \sin(\frac{\pi}{2} - s) \sin t$       **23**  $\theta = \frac{3\pi}{2} +$  multiple of  $2\pi$

**25**  $\theta = \frac{\pi}{4} +$  multiple of  $\pi$       **27** No  $\theta$       **29**  $\phi = \frac{\pi}{4}$       **31**  $|OP| = a, |OQ| = b$

**2**  $\pi, 3\pi, -\frac{\pi}{4}$  radians equal  $180^\circ, 540^\circ, -45^\circ$ . Also  $60^\circ, 90^\circ, 270^\circ$  equal  $\frac{\pi}{3}, \frac{\pi}{2}, \frac{3\pi}{2}$  radians. The alias of  $480^\circ$  is  $120^\circ$  and the alias of  $-1^\circ$  is  $359^\circ$ .

**4**  $\cos 2(\theta + \pi)$  is the same as  $\cos(2\theta + 2\pi)$  which is  $\cos 2\theta$ . Since  $\cos^2 \theta = \frac{1}{2} + \frac{1}{2} \cos 2\theta$ , this also has period  $\pi$ .

**6** Notice the patterns in this table.

**8** Straight distance  $\sqrt{2}$ ; quarter-circle distance  $\frac{\pi}{2}$ ; semicircle distance also  $\frac{\pi}{2}$ .

**10**  $d^2 = (0 - \frac{1}{2})^2 + (1 - \frac{\sqrt{3}}{2})^2 = \frac{1}{4} + 1 - \sqrt{3} + \frac{3}{4} = 2 - \sqrt{3}$ . Then  $12d = 6.21$ . This is the distance around a twelve-sided figure that fits into the circle (curved distance is  $2\pi$ .)

**12** From the inside front cover or the addition formulas:  $\sin(\pi - \theta) = \sin \theta, \cos(\pi - \theta) = -\cos \theta, \sin(\frac{\pi}{2} + \theta) = \cos \theta, \cos(\frac{\pi}{2} + \theta) = -\sin \theta$ .

**14**  $\sin 3t = \sin(2t + t) = \sin 2t \cos t + \cos 2t \sin t$ . This equals  $(2 \sin t \cos t) \cos t + (\cos^2 t - \sin^2 t) \sin t$  or  $3 \sin t - 4 \sin^3 t$ .

**16**  $(\cos t + i \sin t)^2 = \cos^2 t - \sin^2 t + 2i \sin t \cos t$ . Then the double-angle formulas give  $\cos 2t + i \sin 2t$ .

**18** A complete solution is not expected! Finding a point like  $s = \pi/2, t = 3\pi/2$  is not bad.

**20** Formula (9) is  $\sin(s+t) = \sin s \cos t + \cos s \sin t$ . Replacing  $t$  by  $-t$  gives formula (8) for  $\sin(s-t)$ .

(Ask why this replacement is allowed. It is not easy for a student to explain.)

**22**  $\tan(s+t) = \frac{\sin(s+t)}{\cos(s+t)} = \frac{\sin s \cos t + \cos s \sin t}{\cos s \cos t - \sin s \sin t}$ . To simplify, divide top and bottom by  $\cos s$  and  $\cos t$ :  

$$\tan(s+t) = \frac{\tan s + \tan t}{1 - \tan s \tan t}.$$

**24**  $\sec \theta = -2$  when  $\cos \theta = -\frac{1}{2}$ , which happens first at  $\theta = 120^\circ = 2\pi/3$ . Also at  $\theta = 240^\circ = 4\pi/3$ . Then at all angles  $2\pi/3 + 2\pi n$  and  $4\pi/3 + 2\pi n$ .

**26**  $\sin \theta = \theta$  at  $\theta = 0$  and never again. Reason: The right side has slope 1 and the left side has slope  $\cos \theta < 1$ .

(Draw graphs of  $\sin \theta$  and  $\theta$ . A solution with negative  $\theta$  would give a solution for positive  $\theta$  by reversing sign.)

**28**  $\tan \theta = 0$  when  $\theta$  is a multiple of  $\pi$ . The ratio  $y/x$  is zero when  $y = 0$ , so the point on the circle in Figure 1.20 has to be on the  $x$  axis.

**30**  $A \sin(x+\phi)$  equals  $A \sin x \cos \phi + A \cos x \sin \phi$ . Matching with  $a \sin x + b \cos x$  gives  $a = A \cos \phi$  and  $b = A \sin \phi$ . Then  $a^2 + b^2 = A^2 \cos^2 \phi + A^2 \sin^2 \phi = A^2$ . Thus  $A = \sqrt{a^2 + b^2}$  and  $\tan \phi = \frac{A \sin \phi}{A \cos \phi} = \frac{a}{b}$ .

**32** The distance squared from  $(0,0)$  to  $R$  is  $(a + b \cos \theta)^2 + (b \sin \theta)^2$  which simplifies to

$a^2 + 2ab \cos \theta + b^2$ . Notice the parallelogram law:  $(\text{diagonal})^2 + (\text{other diagonal})^2 = 2a^2 + 2b^2$  which is  $(\text{side})^2 + (\text{next side})^2 + (\text{third side})^2 + (\text{fourth side})^2$ .

**34** The amplitude and period of  $2 \sin \pi x$  are both 2.

**36** By Problem 30,  $\sin x + \cos x$  equals  $\sqrt{2} \sin(x + \frac{\pi}{4})$ . The graph should show a sine function with maximum near  $\sqrt{2}$  at  $x = \frac{\pi}{4}$ .

**38** The graph of  $t \sin t$  oscillates between  $\pm 45^\circ$  lines. The graph of  $\sin 4t \sin t$  oscillates inside the graph of  $\sin t$ . See the graph on page 294, at the end of Section 7.2.

MIT OpenCourseWare  
<http://ocw.mit.edu>

Resource: Calculus Online Textbook  
Gilbert Strang

The following may not correspond to a particular course on MIT OpenCourseWare, but has been provided by the author as an individual learning resource.

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.