




Summary: Six Functions, Six Rules, Six Theorems

<i>Integrals</i>	<i>Six Functions</i>	<i>Derivatives</i>
$x^{n+1}/(n+1), n \neq -1$	$x^n$	$nx^{n-1}$
$-\cos x$	$\sin x$	$\cos x$
$\sin x$	$\cos x$	$-\sin x$
$e^{cx}/c$	$e^{cx}$	$ce^{cx}$
$x \ln x - x$	$\ln x$	$1/x$
<b>Ramp function</b>	<b>Step function</b>	<b>Delta function</b>
		

**Six Rules of Differential Calculus**

- The derivative of  $af(x) + bg(x)$  is  $a \frac{df}{dx} + b \frac{dg}{dx}$  **Sum**
  - The derivative of  $f(x)g(x)$  is  $f(x) \frac{dg}{dx} + g(x) \frac{df}{dx}$  **Product**
  - The derivative of  $\frac{f(x)}{g(x)}$  is  $\left( g \frac{df}{dx} - f \frac{dg}{dx} \right) / g^2$  **Quotient**
  - The derivative of  $f(g(x))$  is  $\frac{df}{dy} \frac{dy}{dx}$  where  $y = g(x)$  **Chain**
  - The derivative of  $x = f^{-1}(y)$  is  $\frac{dx}{dy} = \frac{1}{dy/dx}$  **Inverse**
  - When  $f(x) \rightarrow 0$  and  $g(x) \rightarrow 0$  as  $x \rightarrow a$ , what about  $f(x)/g(x)$ ? **L'Hôpital**
- $\lim \frac{f(x)}{g(x)} = \lim \frac{df/dx}{dg/dx}$  if these limits exist. Normally this is  $\frac{f'(a)}{g'(a)}$

**Fundamental Theorem of Calculus**

If  $f(x) = \int_a^x s(t)dt$  then **derivative of integral** =  $\frac{df}{dx} = s(x)$

If  $\frac{df}{dx} = s(x)$  then **integral of derivative** =  $\int_a^b s(x)dx = f(b) - f(a)$

Both parts assume that  $s(x)$  is a continuous function.

**All Values Theorem** Suppose  $f(x)$  is a continuous function for  $a \leq x \leq b$ . Then on that interval,  $f(x)$  reaches its maximum value  $M$  and its minimum  $m$ . And  $f(x)$  takes all values between  $m$  and  $M$  (there are no jumps).

## Summary: Six Functions, Six Rules, Six Theorems

**Mean Value Theorem** If  $f(x)$  has a derivative for  $a \leq x \leq b$  then

$$\frac{f(b) - f(a)}{b - a} = \frac{df}{dx}(c) \text{ at some } c \text{ between } a \text{ and } b$$

“At some moment  $c$ , instant speed = average speed”

**Taylor Series** Match all the derivatives  $f^{(n)} = d^n f / dx^n$  at the basepoint  $x = a$

$$\begin{aligned} f(x) &= f(a) + f'(a)(x-a) + \frac{1}{2}f''(a)(x-a)^2 + \dots \\ &= \sum_{n=0}^{\infty} \frac{1}{n!} f^{(n)}(a) (x-a)^n \end{aligned}$$

Stopping at  $(x-a)^n$  leaves the error  $f^{(n+1)}(c)(x-a)^{n+1}/(n+1)!$

[ $c$  is somewhere between  $a$  and  $x$ ] [ $n=0$  is the Mean Value Theorem]

The Taylor series looks best around  $a=0$   $f(x) = \sum_{n=0}^{\infty} \frac{1}{n!} f^{(n)}(0) x^n$

**Binomial Theorem** shows Pascal's triangle

$$\begin{aligned} (1+x) & \quad \quad \quad \mathbf{1 + 1x} \\ (1+x)^2 & \quad \quad \mathbf{1 + 2x + 1x^2} \\ (1+x)^3 & \quad \mathbf{1 + 3x + 3x^2 + 1x^3} \\ (1+x)^4 & \mathbf{1 + 4x + 6x^2 + 4x^3 + 1x^4} \end{aligned}$$

Those are just the Taylor series for  $f(x) = (1+x)^p$  when  $p = 1, 2, 3, 4$

$$\begin{aligned} f^{(n)}(x) &= (1+x)^p \quad p(1+x)^{p-1} \quad p(p-1)(1+x)^{p-2} \quad \dots \\ f^{(n)}(0) &= \quad \mathbf{1} \quad \quad \quad \mathbf{p} \quad \quad \quad \mathbf{p(p-1)} \quad \dots \end{aligned}$$

Divide by  $n!$  to find the Taylor coefficients = **Binomial coefficients**

$$\frac{1}{n!} f^{(n)}(0) = \frac{p(p-1)\dots(p-n+1)}{n(n-1)\dots(1)} = \frac{p!}{(p-n)!n!} = \binom{p}{n}$$

The series stops at  $x^n$  when  $p = n$  Infinite series for other  $p$

$$\text{Every } (1+x)^p = 1 + px + \frac{p(p-1)}{(2)(1)}x^2 + \frac{p(p-1)(p-2)}{(3)(2)(1)}x^3 + \dots$$

**Practice Questions**

1. Check that the derivative of  $y = x \ln x - x$  is  $dy/dx = \ln x$ .

2. The “sign function” is  $S(x) = \begin{cases} 1 & \text{for } x \geq 0 \\ -1 & \text{for } x < 0 \end{cases}$

What ramp function  $F(x)$  has  $\frac{dF}{dx} = S(x)$ ?  $F$  is the integral of  $S$ .

Why is the derivative  $\frac{dS}{dx} = 2 \delta(x)$ ? (Infinite spike at  $x = 0$  with area 2)

3. (l'Hôpital) What is the limit of  $\frac{2x + 3x^2}{5x + 7x^2}$  as  $x \rightarrow 0$ ? What about  $x \rightarrow \infty$ ?

4. l'Hôpital's Rule says that  $\lim_{x \rightarrow 0} \frac{f(x)}{g(x)} = \frac{f'(0)}{g'(0)}$  when  $f(0) = g(0) = 0$ . Here  $g(x) = x$ .

**5. Derivative is like Difference Integral is like Sum**

Difference of sums If  $f_n = s_1 + s_2 + \dots + s_n$ , what is  $f_n - f_{n-1}$ ?

Sums of differences What is  $(f_1 - f_0) + (f_2 - f_1) + \dots + (f_n - f_{n-1})$ ?

Those are the **Fundamental Theorems** of “**Difference Calculus**”

6. Draw a non-continuous graph for  $0 \leq x \leq 1$  where your function does NOT reach its maximum value.

7. For  $f(x) = x^2$ , which in-between point  $c$  gives  $\frac{f(5) - f(1)}{5 - 1} = \frac{df}{dx}(c)$ ?

8. If your average speed on the Mass Pike is 75, then at some instant your speedometer will read \_\_\_\_\_.

9. Find three Taylor coefficients  $A, B, C$  for  $\sqrt{1+x}$  (around  $x = 0$ ).

$$(1+x)^{\frac{1}{2}} = A + Bx + Cx^2 + \dots$$

10. Find the Taylor (= Binomial) series for  $f = \frac{1}{1+x}$  around  $x = 0$  ( $p = -1$ ).

MIT OpenCourseWare  
<http://ocw.mit.edu>

Resource: Highlights of Calculus  
Gilbert Strang

The following may not correspond to a particular course on MIT OpenCourseWare, but has been provided by the author as an individual learning resource.

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.